

RESEARCH

Open Access

Prediction of T-cell epitopes of hepatitis C virus genotype 5a

Maemu P Gedezha, M Jeffrey Mphahlele and Selokela G Selabe*

Abstract

Background: Hepatitis C virus (HCV) is a public health problem with almost 185 million people estimated to be infected worldwide and is one of the leading causes of hepatocellular carcinoma. Currently, there is no vaccine for HCV infection and the current treatment does not clear the infection in all patients. Because of the high diversity of HCV, protective vaccines will have to overcome significant viral antigenic diversities. The objective of this study was to predict T-cell epitopes from HCV genotype 5a sequences.

Methods: HCV near full-length protein sequences were analyzed to predict T-cell epitopes that bind human leukocyte antigen (HLA) class I and HLA class II in HCV genotype 5a using Propred I and Propred, respectively. The Antigenicity score of all the predicted epitopes were analysed using VaxiJen v2.0. All antigenic predicted epitopes were analysed for conservation using the IEDB database in comparison with 406, 221, 98, 33, 45, 45 randomly selected sequences from each of the HCV genotypes 1a, 1b, 2, 3, 4 and 6 respectively, downloaded from the GenBank. For epitope prediction binding to common HLA alleles found in South Africa, the IEDB epitope analysis tool was used.

Results: A total of 24 and 77 antigenic epitopes that bind HLA class I and HLA class II respectively were predicted. The highest number of HLA class I binding epitopes were predicted within the NS3 (63%), followed by NS5B (21%). For the HLA class II, the highest number of epitopes were predicted in the NS3 (30%) followed by the NS4B (23%) proteins. For conservation analysis, 8 and 31 predicted epitopes were conserved in different genotypes for HLA class I and HLA class II alleles respectively. Several epitopes bind with high affinity for both HLA class I alleles and HLA class II common in South Africa.

Conclusion: The predicted conserved T-cell epitopes analysed in this study will contribute towards the future design of HCV vaccine candidates which will avoid variation in genotypes, which in turn will be capable of inducing broad HCV specific immune responses.

Keywords: Hepatitis C virus, Genotype 5a, *In-silico*, T-cell epitopes, Vaccines

Introduction

Hepatitis C virus (HCV) is estimated to infect 185 million people worldwide [1]. Chronic HCV infection leads to progressive liver disease, being one of the major causes of hepatocellular carcinoma and one of the most common indications for liver transplantation [2]. The World Health Organization (WHO) strongly recommends combination therapy with pegylated interferon and ribavirin for chronically infected patients who qualify for treatment

[1]. Recently, two NS3 protease inhibitors (boceprevir and telaprevir) have been approved by the US Food and Drug Administration, and the WHO conditionally (until more evidence has accumulated) recommends that these drugs should be given in combination with pegylated interferon and ribavirin for the treatment of chronic HCV genotype 1 infections. Also, the WHO strongly recommends that sofosbuvir be given in combination with ribavirin alone in patients who cannot tolerate interferon and are chronically infected with genotypes 1, 2, 3 and 4 [1]. However, these therapies are still not affordable in most developing countries.

* Correspondence: selokela.selabe@ul.ac.za

HIV and Hepatitis Research Unit, Department of Virology, University of Limpopo, Medunsa Campus/National Health Laboratory Service, Pretoria, South Africa

As a result, the development of an effective HCV vaccine is undoubtedly the best solution for the ultimate control of HCV infections, and is a public health priority.

Prophylactic vaccines against viral infections are generally aimed at inducing a humoral (B-cell) immune response, while therapeutic vaccines preferably activate both humoral and cellular (T-cell) immune responses [3]. A successful HCV vaccine will need to stimulate both arms of the adaptive immune response, since while both cellular and humoral immune responses occur in a naturally infected host, the current consensus is that a strong cellular response is vital for viral clearance and protection [4].

The development of an effective HCV vaccine requires an understanding of the host's adaptive immune response to natural infection. As with other viral infections, viral antigens are presented to CD4+ and CD8+ T-cells via human leukocyte antigen (HLA) class II and class I molecules, respectively [5]. Different HLA class alleles have been found to be associated with HCV infection. For example, HLA-A*11, HLA-Cw*04 and HLA-B*53 have been associated with HCV persistence [6,7], while HLA-B*27, HLA-A*11:01, HLA-B*57, HLA-Cw*01:02 and HLA-A*03 have been associated with spontaneous HCV clearance [8,9]. Also, HLA-DRB1*11 and HLA-DQB1*03:01 have been associated with decreased disease severity of HCV infection globally, suggesting they may present HCV-derived epitopes more efficiently to CD4+ T-cells than others and thus capable of viral clearance [10]. During acute HCV infection, development and persistence of strong specific responses by CD8+ and CD4+ T-cells [11,12] and neutralizing antibodies [13] are associated with viral clearance, with HCV-specific CD8+ and CD4+ T-cells usually being transient or absent in patients who develop persistent infections. The rate of chronic liver disease progression has been shown to be determined by the magnitude of HCV-specific CD4+ T-cell responses, since these cells are essential for both the cellular and humoral responses [14]. CD8+ T-cells are essential for long-term protection against chronic HCV [15], while CD4+ T-cells play a role in viral clearance [16].

HCV infection evades the host's immune system by generating immune escape variants through alteration of the virus HLA-restricted epitopes to avoid being recognized by T-cells and neutralizing antibodies [14]. Thus effective HCV vaccines will need to target protective epitopes that display minimal cross-genotype amino acid variability as they will provide broad potency [17]. Peptides corresponding to protective epitopes are desirable vaccine candidates because they are easy to construct and produce, and they do not contain infectious materials [18]. The first step in the process

of epitope-based vaccine design and development is the *in-silico* prediction of peptide binding affinities to HLA proteins [19].

Genotype 5 accounts for over 50% of HCV infections in South Africa [20], and is becoming more prevalent in Europe and North America [21]. It is the most conserved of HCV genotypes, being classified into only one subtype (5a) [22-24]. The growing prevalence of HCV genotype 5a in different parts of the world necessitates its molecular characterization in order to improve the formulations of vaccine candidates that are in development. Thus the aim of this study was to assess immunological determinants by predicting conserved epitopes in near-full length HCV genotype 5a sequences using a suite of online programmes to help in the designing of new vaccine candidates.

Results

Prediction of T-cell epitopes

For HLA class I, a total of 24 antigenic epitopes were predicted in the consensus near full-length of genotype 5a (Table 1). Epitope NS3¹³²⁵⁻¹³³³ covered 30 of the 47 HLA class I alleles that were analysed assuring high binding affinity to different alleles. For conservation analysis with other genotypes, 8 of 24 epitopes were 100% conserved for specific genotypes. Epitopes NS3¹³³²⁻¹³⁴⁰ and NS5B²⁵⁵⁷⁻²⁵⁶⁵ were highly conserved in all genotypes analysed, and epitope E2⁶⁸⁴⁻⁶⁹² was conserved in all genotypes except for genotype 2, while 5 other epitopes were conserved in either 2 or 3 genotypes analysed. In addition epitope NS4B¹⁸³²⁻¹⁸⁴⁰ was conserved in genotype 1a, 1b, 2 and 4 epitope variants at anchor residues position 2 and 9 while E2⁶⁷⁷⁻⁶⁸⁵, NS3¹³²⁵⁻¹³³³ and NS3¹³⁵⁷⁻¹³⁶⁵ were conserved in at least 3 genotypes epitope variants each (Table 1). For HLA class II, 77 epitopes were predicted (Table 2). Epitope NS4B¹⁸⁷⁹⁻¹⁸⁸⁷ and NS4B¹⁸⁸⁰⁻¹⁸⁸⁸ covered 51 of 51 HLA class II alleles analysed. For conservation analysis with other genotypes, 31 of 71 epitopes were conserved for specific genotypes. Some epitopes were highly conserved in all genotypes (E2⁵⁰⁷⁻⁵¹⁵, E2⁵⁰⁹⁻⁵¹⁷, NS3¹²⁵³⁻¹²⁶¹, NS3¹²⁵⁴⁻¹²⁶², NS3¹³²⁷⁻¹³³⁵, NS3¹³⁹²⁻¹⁴⁰⁰, NS4B¹⁹¹⁶⁻¹⁹²⁴ and NS4B¹⁹¹⁹⁻¹⁹²⁷), while other epitopes were conserved in at least 1 to 4 of the genotypes analysed. Epitope NS4B¹⁸⁸⁶⁻¹⁸⁹⁴ was conserved at anchor residues position 1, 4, 6 and 9 in all 6 genotypes epitope variants while E2⁶⁹²⁻⁷⁰⁰, NS2⁹⁶⁴⁻⁹⁷², NS3¹⁴¹⁸⁻¹⁴²⁶ and NS4¹⁵⁶¹⁻¹⁵⁶⁹ were conserved in at least 5 genotypes epitope variants each (Table 2). Epitopes NS3¹⁵⁸⁵⁻¹⁵⁹³, NS5A²²⁸⁵⁻²²⁹³ and NS5B²⁸⁸⁹⁻²⁸⁹⁷ were predicted to cover both HLA class I and HLA class II alleles. The highest number of HLA class I binding epitopes were predicted within the NS3 (63%), followed by NS5B (21%), and for the HLA class II, the highest number of

Table 1 HLA class I predicted epitopes of HCV genotype 5a and their antigenicity prediction score, number of allele and conservation (in percentages) in different genotypes

Position	Epitope sequence	No of allele	Antigenicity score	Genotype 1a	Genotype 1b	Genotype 2	Genotype 3	Genotype 4	Genotype 6
E2									
651	RCDLEDRDR	9	2.7428						
677	CSFTTPAL	27	0.8772	CSFT7LPAL (94)	CSFT7LPAL (92)		CSFTMPAL (73)		
684	ALSTGLIHL [#] [25-27]	29	0.9005	ALSTGLIHL (70)	ALSTGLIHL (99)	ALSTGLIHL (97)	ALSTGLIHL (97)	ALSTGLIHL (84)	ALSTGLIHL (91)
NS2									
861	VPPLQVRGG	6	1.4312	VPPLMVRGG (85)					
NS3									
1032	YAQQTRGVL	26	0.7638	YAQQTRGLL (99)					
1033	AQQTRGVLG	5	0.7751	AQQTRGLLG (99)					
1128	ADLYLVTRH	5	1.0277	SDLYLVTRH (99)	SDLYLVTRH (99)	VDLYLVTRN (92)			
1325	TILGIGTVL	30	0.7555	SILGIGTVL (93)		TILGIGTVL (88)	SILGIGTVL (94)	TILGIGTVL (82)	
1332	VLDQAETAG	5	0.5004	VLDQAETAG (96)	VLDQAETAG (98)	VLDQAETAG (96)	VLDQAETAG (100)	VLDQAETAG (93)	VLDQAETAG (71)
1357	TPHPNIEEV	24	0.8423	VPHNIEEV (89)	VPHNIEEV (80)		VPHSNIEEV (94)		
1359	HPNIEEVAL [#] [28-31]	24	1.1654	HPNIEEVAL (88)	HPNIEEVAL (76)		HSNIEEVAL (94)		
1370	EGEIPFYGR	9	1.0560	TGEIPFYGK (97)					
1374	PFYGRAIPL	7	0.9479	PFYGKAIPL (99)	PFYGKAIPL (85)	PFYGKAIPL (82)			PFYGKAIPL (73)
1642	TKYIMACMS	6	0.7008	PFYGKAIPL (99)	TKYIMACMS (70)		TKYIMACMS (70)	TKYIMACMS (93)	TKYIMTCMS (87)
NS4B									
1712	SASLPYMDE	5	0.5488						
1819	QIAPPTAAT	5	0.5293	QLAAPGAAT (99)	QLAPPSAAS (94)	QIAPPAGAT (91)		QIATPTAST (89)	
1832	SGMAGAAVG	5	0.8750	AGLAGAAIG (81)	AGIAGAAVG (93)	SGLVGA AVG (98)	SGLAGAAIG (76)		
1848	LIDILAGYG	5	0.8431	LVDILAGYG (94)	LVDILAGYG (95)		LLDILAGYG (79)	LVDILAGYG (84)	
1854	GYGAGVAGA	9	0.5562	GYGAGVAGA (99)	GYGAGVAGA (99)		GYGAGVSGA (94)	GYGAGVAGA (91)	GYGAGVSGA (84)
NS5B									
2522	GYGAKEVRS	7	1.3540	GYGAKDVRC (95)	GYGAKDV RN (84)	GFGAKEVRS (93)	GYSAKDVRS (79)		
2557	TIMAKNEVF [#] [32]	12	0.6303	TIMAKNEVF (99)	TIMAKNEVF (80)	TIMAKNEVF (83)	TIMAKNEVF (100)	TIMAKNEVF (84)	TIMAKNEVF (93)
2568	EPSKGGKKP	6	1.4218	QPEKGG RKP (93)					
2720	LASCRAAKL	28	0.5264						
2886	LHGLSAFSL	11	0.6732	LHGLSAFSL (99)	LHGLSAFSL (98)			LHGLSAFTL(82)	LHGMAAFSL (98)

[#]-indicates that the epitope has been experimentally proven to be a true positive.

Bold- indicates percentage of epitope that is 100% conserved in more than 70% of the sequences analysed in each genotype.

Italic- indicates amino acid(s) variation in epitope in comparison to the predicted epitope.

Table 2 HLA class II predicted epitopes of HCV genotype 5a and their antigenicity prediction score, number of allele and conservation (in percentages) in different genotypes

Position	Epitope sequence	No of allele	Antigenicity score	Genotype 1a	Genotype 1b	Genotype 2	Genotype 3	Genotype 4	Genotype 6
E1									
320	WMMMMNWSP	10	1.3610	WMMMMNWSP (99)	WMMMMNWSP (97)		WMMMMNWSP (97)	WMMMMNWSP (93)	WMMMMNWSP (93)
E2									
507	YCFTSPVV	6	1.2112	YCFTSPVV (93)	YCFTSPVV (98)	YCFTSPVV (88)	YCFTSPVV (100)	YCFTSPVV (91)	YCFTSPVV (91)
509	FTPSPVVVG	17	1.3553	FTPSPVVVG (94)	FTPSPVVVG (98)	FTPSPVVVG (88)	FTPSPVVVG (100)	FTPSPVVVG (89)	FTPSPVVVG (73)
665	LLHTTTQWA	5	0.6616	LLSTTQWQ (100)			LLHSTTELA (73)	LLSTTQWQ (73)	
666	LHTTTQWAI	25	0.5863	LTTTQWQI (100)			LHSTTELA (73)		
692	LHQNIVDTQ	6	0.8564	LHQNIVDVQ (100)	LHQNIVDVQ (73)	LHQNIVDVQ (95)	LHQNIVDVQ (100)	LHQNIVDVQ (87)	
739	LLVCAEAA	22	0.5506	LLISQAEAA (100)	LLIAQAEAA (72)		LMISQAEAA (73)		
P7									
803	LPHRALLD	7	0.6966	LPQRAYALD (100)	LPPRAYAMD (86)				
NS2									
860	WVPLQVRG	7	1.1998	WVPLMVRG (100)			WVPLLARG (82)		
861	VPPLQVRGG	25	1.4312	VPPLMRGG (100)					
864	LQVRGGRDA	20	0.6368	LWVRGGRDA (100)	LWVRGGRDA (86)				
880	FHPALGFEI	19	1.4374	VHPALVFDI (99)					
892	LLGILGPLY	14	0.5940	LLAVLGPLW (100)			LIAVLGPLY (82)		
895	ILGPLYLLQ	5	0.8264	VLGPLWLLQ (99)			VLGPLYLIQ (88)		
938	LLHLGRLTG	25	0.5554						
964	LRDLAVATE	9	1.0497	LRDLAVAVE (100)	LRDLAVAVE (93)	LRDLAVAVE (85)	LKDLAVATE (73)		LRDLAVAVE (93)
997	LAGLPVSAR	11	0.9120	LAGLPVSAR (100)			LCGLPVSAR (100)		
1025	LLAPITAYA	22	0.5253	LLAPITAYA (99)			LLAPITAYA (79)	LLAPITAYA (89)	
NS3									
1047	LTGRDKNEA	22	1.5633	LTGRDKNQV (100)	LTGRDKNQV (93)		LTGRDKNV (90)		
1131	YLVTRHADV [#] [33-35]	15	0.7569	YLVTRHADV (99)	YLVTRHADV (99)	YLVTRNADV (97)			
1152	LLSPRPISY	5	2.2999	LLSPRPISY (99)	LLSPRPVSY (73)	LLSPRPLST (89)			
1153	LSPRPISYL	5	1.3109	LSPRPISYL (99)	LSPRPVSYL (73)	LSPRPLSTL (89)			
1253	VLNPSVAAT	14	1.2489	VLNPSVAAT (100)	VLNPSVAAT (100)	VLNPSVAAT (97)	VLNPSVAAT (100)	VLNPSVAAT (96)	VLNPSVAAT (89)
1254	LNPSVAATL	6	0.9169	LNPSVAATL (100)	LNPSVAATL (100)	LNPSVAATL (98)	LNPSVAATL (100)	LNPSVAATL (95)	LNPSVAATL (89)
1258	VAATLGFGA	11	1.2617	VAATLGFGA (90)	VAATLGFGA (85)	VAATLGFGA (98)			
1262	LGFGAYMSK [#] [36-38]	14	0.8784	LGFGAYMSK (87)	LGFGAYMSK (82)	LGFGAYMSK (78)			

Table 2 HLA class II predicted epitopes of HCV genotype 5a and their antigenicity prediction score, number of allele and conservation (in percentages) in different genotypes (Continued)

1264	FGAYMSKAY	14	0.6561	FGAYMSKAH (100)	FGAYMSKAH (82)					
1327	LGIGTVLDQ	42	0.5607	LGIGTVLDQ (97)	LGIGTVLDQ (88)	LGIGTVLDQ (98)	LGIGTVLDQ (100)	LGIGTVLDQ (93)	LGIGTVLDQ (96)	
1373	IPFYGRAIP	16	0.5679	IPFYGKAIP (100)	IPFYGKAIP (93)	IPFYGKAIP (81)				
1375	FYGRAIPLA	22	1.0280	FYGKAIPLE (100)	FYGKAIPLE (81)					FYGKAIPLE (71)
1392	IFCHSKKKC	22	1.5437	IFCHSKKKC (93)	IFCHSKKKC (94)	IFCHSKKKC (95)	IFCHSKKKC (88)	IFCHSKKKC (80)	IFCHSKKKC (93)	
1415	VAYYRGLDV	21	0.8250	VAYYRGLDV (98)	VAYYRGLDV (96)	VAYYRGLDV (92)	VAYYRGLDV (79)	VAYYRGLDV (91)	VAYYRGLDV (91)	VAFYRGVDV (89)
1418	YRGLDVAVI	51	1.0522	YRGLDVSVI (100)	YRGLDVSVI (97)	YRGLDVSVI (84)	YRGLDVSVI (85)	YRGLDVSVI (91)	YRGLDVSVI (91)	YRGLDVSVI (67)
1482	VSRQRGR	12	2.0649	VSRQRGR (100)	VSRQRGR (93)	VSRQRGR (96)	VSRQRGR (97)	VSRQRGR (98)	VSRQRGR (98)	VSRQRGR (73)
1561	VFTGLTNID	9	1.1689	VFTGLTHID (100)	VFTGLTHID (96)	VFTGLTHID (98)	VFTGLTHID (97)			VFTGLTHID (89)
1583	FPYLVAYQA	24	0.6584		FPYLVAYQA (85)					FAYLVAYQA (78)
1585	YLVAQATV[#] [25,39]	17	0.5020	YLVAQATV (99)	YLVAQATV (85)	YLTAYQATV (70)		YLVAQATV (87)	YLVAQATV (91)	
1586	LVAYQATVC	28	0.6052	LVAYQATVC (99)	LVAYQATVC (85)			LVAYQATVC (82)	LVAYQATVC (91)	
1631	VQNEITLTH	35	1.1979	VQNEITLTH (100)				VQNEITLTH (80)		
1641	ITKYIMACM	8	0.6964	ITKYIMTCM (100)		ITKYIATCM (94)		ITKYIMACM (75)		ITKYIMTCM (87)
1645	IMACMSADL	22	0.8406	IMTCMSADL (100)	IMACMSADL (82)	IATCMQADL (99)	IMACMSADL (70)	IMACMSADL (93)		IMTCMSADL (87)
NS4B										
1733	LGLIGTAGQ	43	1.2450							
1765	MWNFVSGIQ	18	1.2300	MWNFVSGIQ (100)	MWNFVSGIQ (95)	MWNFVSGIQ (96)	MWNFVSGIQ (100)	MWNFVSGIQ (87)		
1766	WNFVSGIQY	22	0.6246	WNFVSGIQY (100)	WNFVSGIQY (95)	WNFVSGIQY (96)	WNFVSGIQY (100)	WNFVSGIQY (87)		WNFVSGIQY (91)
1791	MSFTAAVTS	8	0.7414	MAFTAAVTS (100)	MAFTASITS (88)	MAFSAALTS (93)	MAFTASVTS (92)	MSFTAAVTS (93)		
1812	LGGWVASQI	16	0.608	LGGWVAAQL (100)	LGGWVAAQL (95)		LGGWVATHL (88)	LGGWVASQI (96)		
1849	IDILAGYGA	9	0.7525	VDILAGYGA (100)	VDILAGYGA (97)		LDILAGYGA (79)	VDILAGYGA (84)		
1863	LVAFKIMCG	47	1.0160	LVAFKIMSG (100)		LVAFKIMSG (95)	LVAFKIMGG (100)	WTFKIMSG (89)		LVAFKIMSG (84)
1879	LVNLLPSIL	51	0.7592	LVNLLPAIL (100)	LVNLLPAIL (90)		MVNLLPAIL (92)	LVNLLPAIL (73)		
1880	VNLLPSILC	51	0.9874	VNLLPAILS (100)	VNLLPAILS (92)		VNLLPAILS (94)	VNLLPAILS (89)		VNLLPAILS (82)
1883	LPSILCPGA	6	0.7104	LPAILSPGA (100)	LPAILSPGA (99)	LPAILSPGA (74)	LPAILSPGA (100)	LPAILSPGA (89)		LPAILSPGA (84)
1886	ILCPGALVV	9	0.9365	ILSPGALW (100)	ILSPGALW (99)	ILSPGALW (98)	ILSPGALW (100)	ILSPGALW (91)		ILSPGALW (89)
1892	LVGVVCAA	30	0.7097	LWGVVCAA (100)	LWGVVCAA (98)	LVGVVCAA (97)	LVGVVCAA (94)	LWGVVCAA (89)		LWGVVCAA (93)
1893	VGVVCAA	50	0.9530	VGVVCAA/ (100)	VGVVCAA/ (96)	VGVVCAA/ (98)	VGVVCAA/ (85)	VGVVCAA/ (91)		VGVVCAA/ (76)
1896	VCAA/LRR	35	0.9887	VCAA/LRR (100)	VCAA/LRR (95)	VCAA/LRR (97)	VCAA/LRR (85)	VCAA/LRR (91)		VCAA/LRR (76)
1897	ICAA/LRRH	12	1.2316	VCAA/LRRH (100)	VCAA/LRRH (97)	ICAA/LRRH (97)	ICAA/LRRH (85)	VCAA/LRRH (89)		
1916	MNRLIAFAS	48	0.5291	MNRLIAFAS (100)	MNRLIAFAS (99)	MNRLIAFAS (99)	MNRLIAFAS (100)	MNRLIAFAS (93)		MNRLIAFAS (96)

Table 2 HLA class II predicted epitopes of HCV genotype 5a and their antigenicity prediction score, number of allele and conservation (in percentages) in different genotypes (Continued)

1919	LIAFASRGN	24	2.0235	LIAFASRGN (100)	LIAFASRGN (99)	LIAFASRGN (98)	LIAFASRGN (100)	LIAFASRGN (93)	LIAFASRGN (96)
1922	FASRGNHVS	18	1.2938	FASRGNHVS (100)	FASRGNHVS (98)	FASRGNHVA (97)	FASRGNHVS (94)	FASRGNHVS (82)	FASRGNHVS (93)
NS5A									
1994	WLQAKLLPQ	33	1.1603	WLKAKLMPQ (100)	WLQSKLLPR (84)				
2101	YITGVTQDN	6	1.1358	YVTGMTTDN (100)					
2105	VTQDNLKCP	6	1.0178	MTTDNLKCP (100)					
2237	MGGNITRVE	20	1.2656	MGGNITRVE (92)	MGGNITRVE (96)		MGSNITRVE (91)		
2285	LPVWARPGY	9	0.6143	LPWARPDY (100)					LPWARPDY (78)
NS5B									
2422	MSYSWTGAL	9	0.5749	MSYSWTGAL (76)	MSYWTGAL (94)	MSYSWTGAL (97)	MSYSWTGAL (85)	MSYSWTGAL (91)	
2451	LRHHNLVYS	40	1.2321	LRHHNLVYS (89)	LRHHNMVYA (86)		LRHHNLVYS (97)		
2523	YGAKEVRSL	24	1.1739	YGAKDVRCH (100)	YGAKDVRNL (85)	FGAKEVRSL (92)	YSAKDVRSL (76)		
2559	MAKNEVFAV	18	0.8466	MAKNEVFCV (100)	MAKNEVFCV (81)	MAKNEVFCV (65)	MAKNEVFCV (85)		MAKNEVFCV (96)
2564	VFAVEPSKG	10	0.7925	VFCVQPEKG (100)	VFCVQPEKG (89)				
2637	FSYDTRCFD	11	1.5808	FSYDTRCFD (99)		FSYDTRCFD (95)	FSYDTRCFD (100)	FSYDTRCFD (93)	FSYDTRCFD (87)
2660	YQSCDLQPE	11	1.3802	YQCCDLDPQ (100)	YQCCDLAPE (94)				
2696	YRRCRASGV	41	1.2642	YRRCRASGV (99)	YRRCRASGV (99)	YRRCRASGV (98)	YRRCRASGV (97)		
2778	YDLELVTS	6	1.0923	YDLELVTS (100)	YDLELVTS (96)	YDLELVTS (98)	YDLELVTS (91)		YDLELVTS (96)
2850	FSVLQSSEQ	21	0.8451		FSLLAQEQ (84)			FSLQSSEA (78)	
2871	VYSVTPLDL	29	2.0113	CYSIEPLDL (100)	CYSIEPLDL (75)		TYSVTPLDL (91)	TYSVTPLDL (87)	
2883	IQLHGLSA	50	0.4322	IQLHGLSA (98)	IQLHGLSA (72)	IQLAGLST (93)	IQLHGLSA (79)	IQLHGLSA (75)	IQLHGLMAA (96)
2889	LSAFSLHSY [#] [40]	6	0.6007	LSAFSLHSY (99)	LSAFSLHSY (97)			LSAFSLHGY (80)	MAAFSLHGY (89)

[#]-indicates that the epitope has been experimentally proven to be a true positive.

Bold- indicates percentage of epitope that is 100% conserved in more than 70% of the sequences analysed in each genotype.

Italic- indicates amino acid(s) variation in epitope in comparison to the predicted epitope.

epitopes were predicted in the NS3 (30%) followed by the NS4B (23%) proteins (Table 3).

Epitope binding affinity to common South African HLA class I and HLA class II alleles

Genotype 5 epitopes and their genotypic variants were analysed for their binding affinity to HLA class I and HLA class II alleles most common in South Africa, where genotype 5a is predominating. For HLA class I, 11 of the most common South African HLA-A alleles (HLA-A*01:01, HLA-A*02:01, HLA-A*30:01 and HLA-A*30:02), HLA-B (HLA-B*0702, HLA-B*08:01 and HLA-B*3501), and HLA-C (HLA-C*04:01, HLA-C*06:01, HLA-C*07:01 and HLA-C*07:02) were analysed. The limitation of ProPred 1 is that it does not cover most of the main HLA class I alleles: HLA-A (HLA-A*01:01, HLA-A*30:01 and HLA-A*30:02), HLA-B (HLA-B*08:01) and HLA-C (HLA-C*04:01, HLA-C*06:01, HLA-C*07:01 and HLA-C*07:02) that are observed in South Africa. As a result the IEDB epitope analysis tool was used to predict epitopes of all the 11 most common HLA-A, HLA-B, HLA-C covering HLA class I alleles found in South Africa with the ANN prediction server. Thirteen antigenic epitopes with high binding affinity score of <50 IC₅₀nM were predicted. Most epitopes bind with high affinity to single HLA class I alleles with exception of epitope NS5B²⁸⁸⁹⁻²²⁹⁷ LSAFSLHSY that bind with high affinity to 2 HLA-A alleles (HLA-A*01:01 and HLA-A*30:02). Four of the epitopes binding to HLA-B*35:01 followed by HLA-A*02:01 with 3 epitopes binding to it, however, none of the epitopes bind to HLA-C*04:01, HLA-C*06:01 and HLA-C*07:02 alleles. The NS3¹³⁵⁹⁻¹³⁶⁷ shows a level of promiscuity to HLA-A and HLA-B alleles. NS3¹³⁵⁹⁻¹³⁶⁷ HPNIEEVAL bind with intermediate affinity to HLA-B*07:02, while its genotype 2, 3 and 4 variant HSNIEEVAL bind with poor affinity and genotype 6 variant HPNITETAL bind with high affinity. For HLA-B*35:01, the NS3¹³⁵⁹⁻¹³⁶⁷ HPNIEEVAL and genotype 6 variant HPNITETAL bind with high affinity while genotype 2, 3 and 4 variant HSNIEEVAL bind with poor affinity. Three (E2⁶⁸⁴⁻⁶⁹², NS3¹⁰³²⁻¹⁰⁴⁰ and NS3¹³⁵⁹⁻¹³⁶⁷) of the thirteen epitopes were predicted by both ProPred 1 and IEDB analysis tool (Table 4).

For HLA class II alleles, 4 most common HLA-DRB alleles (HLA-DRB1*03:01, HLA-DRB1*04:01, HLA-DRB1*11:01 and HLA-DRB1*15:01) were analysed. Nineteen antigenic

epitopes with high binding affinity score of <50 IC₅₀nM were predicted. The HLA-DRB1*11:01 has the highest number of binding epitopes (10) followed by HLA-DRB1*04:01 and HLA-DRB1*15:01 (7 each). Epitopes NS4B¹⁹¹⁹⁻¹⁹²⁷ LIAFASRGN was predicted to be the most promiscuous epitope binding to HLA-DRB1*04:01, HLA-DRB1*11:01 and HLA-DRB1*15:01 with high affinity and HLA-DRB1*03:01 with intermediate affinity. This epitope is conserved in all genotypes. Epitopes E2⁵⁰⁷⁻⁵¹⁵, NS4^{B1774-1782}, NS4B¹⁹¹⁹⁻¹⁹²⁷, and NS4B¹⁹²⁰⁻¹⁹²⁸ were highly conserved in all genotypes (Table 5).

Validation of epitopes

Seven of the predicted epitopes were previously confirmed experimentally by other studies as true positives in comparison with the epitopes analysed in the IEDB resource database. Majority of the epitopes predicted in this study have not been previously tested experimentally. The 'true epitopes' are highlighted by (#) in Tables 1, 2 and 4.

Discussion

Several studies that have published HCV epitopes focused mainly on genotype 1 [41,42], but most of these studies do not take into account the diversity in other genotypes that are common in developing countries like most African countries. In the present study, predicted antigenic epitopes of HCV genotype 5a proteins from South Africa were analysed followed by conservation with randomly selected genotypes 1-6 references from GenBank. Several studies have confirmed the importance of using immunoinformatics as good predictors for selecting HLA ligands, T-cell epitopes and immunogenicity [43]. As a result, several immunoinformatics methods have been developed to assist in the identification of HLA binding peptides [44,45].

For this analysis, near full-length sequences covering all HCV proteins with the exclusion of the 3' end of the NS5B were included to maximize number of epitopes predicted. The use of the whole viral genome for developing epitope vaccines has a potential control over the immune response and eliminating the side effects [43], and it also increases the chance of detecting a virus at any developmental stage [46]. It has been shown that multiple epitopes from different parts of the HCV genome are important to produce a vaccine that can elicit strong humoral immune responses and multiple specific cellular

Table 3 Distribution of genotype 5a HLA class I and II predicted epitopes in each of the HCV gene

HLA class	No of predicted epitopes	Distribution of the predicted epitopes in each of the HCV gene ^a									
		C	E1	E2	P7	NS2	NS3 ^b	NS4A	NS4B	NS5A	NS5B
I	24	0 (0)	0 (0)	3 (12)	0 (0)	1 (4)	15 (63)	0 (0)	0 (0)	0 (0)	5 (21)
II	77	0 (0)	1 (1)	6 (8)	1 (1)	10 (13)	23 (30)	0 (0)	18 (23)	5 (7)	13 (7)

^aNumber of epitopes and their percentage in brackets.

^bHCV gene with highest number of binding epitopes (in bold).

Table 4 Binding affinity scores of predicted epitopes and their variants to common HLA I allele types prevalent in South Africa

Gene	Epitope sequence ^a	Genotype of epitope	HLA I allele types ^b										
			HLA-A				HLA-B			HLA-C			
			HLA-A*01:01	HLA-A*02:01	HLA-A*30:01	HLA-A*30:02	HLA-B*07:02	HLA-B*08:01	HLA-B*35:01	HLA-C*04:01	HLA-C*06:02	HLA-C*07:01	HLA-C*07:02
E2													
684	ALSTGLIHL [#] [25-27]	1a, 1b, 3, 4,5a, 6	+	+++	+	+	+	+	+	+	+	+	+
	ALSTGLLHL	2	+	+++	+	+	+	+	+	+	+	+	+
NS2													
1025	LLAPITAYA	1a, 3, 4, 5a	+	+++	+	++	+	+	+	+	+	+	+
	LLAPITAYT	2	+	+++	+	++	+	+	+	+	+	+	+
NS3													
1032	YAQQTRGVL	5a	+	+	+	+	+++	+	+	+	+	+	+
	YAQQTRGLL	1a, 2	+	+	+	+	++	+	+	+	+	+	+
	YSQQTRGLL	1b	+	+	+	+	+	+	+	+	+	+	+
	YAQQTRGLV	6	+	+	+	+	+	+	+	+	+	+	+
1242	AAAYAAQGYK	1a, 1b, 4, 5a	+	+	+++	+	+	+	+	+	+	+	+
	AAAYASQGYK	2	+	+	+++	+	+	+	+	+	+	+	+
1264	FGAYMSKAY	5a	+	+	+	+	+	+	+	+++	+	+	+
	FGAYMSKAH	1a, 1b, 2	+	+	+	+	+	+	+	+	+	+	+
1359	HPNIEEVAL [#] [28-31]	1a, 1b, 5a	+	+	+	+	++	+	+	+++	+	+	+
	HSNIEEVAL	2, 3, 4	+	+	+	+	+	+	+	+	+	+	+
	HPNITEVAL	6	+	+	+	+	+++	+	+	+++	+	+	+
1367	LPSEGEIPF	5a	+	+	+	+	++			+++	+	+	+
	LGHEGEIPF	2	+	+	+	+	+	+	+	++	+	+	+
	LGSEGEIPF	3	+	+	+	+	+	+	+	++	+	+	+
	LPTTGEIPF	4, 6	+	+	+	+	++			+++	+	+	+
1585	YLVAYQATV [#] [25,39]	1a, 1b, 4, 5a, 6	+	+++	+	+	+	+	+	+	+	+	+
	YLTAYQATV	2, 3	+	+++	+	+	+	+	+	+	+	+	+
NS5A													
1927	NHVSPHYV	1a, 1b, 3, 4,5a, 6	+	+	+	+	+	+	+	+	++	+++	+
	NHVAPHYV	2	+	+	+	+	+	+	+	+	+	++	+
2285	LPWARPQY	5a	+	+	+	+	+	+	+	+++	+	+	+
	LPWARPQY	1a, 3, 4, 6	+	+	+	+	+	+	+	+++	+	+	+
	LPAWARPQY	2	+	+	+	+	+	+	+	+++	+	+	+

Table 4 Binding affinity scores of predicted epitopes and their variants to common HLA I allele types prevalent in South Africa (Continued)

NS5B													
2696	YRRCRASGV	1a, 1b, 2, 3, 5a	+	+	+	+	+	++	+	+	++	+++	++
2763	MTRYSAAPP	1a, 1b, 2, 3, 4, 5a, 6	+	+	+++	+	+	+	+	+	+	+	+
2889	LSAFSLHSY [#] [40]	1a, 1b, 5a	+++	+	+	+++	+	+	++	+	+	+	+
	LSAF7LHSY	3	+++	+	+	+++	+	+	++	+	+	+	+
	LSAF7LHGY	4	+++	+	+	+++	+	+	+	+	+	+	+

[#]-indicates that the epitope has been experimentally proven to be a true positive.

^a-italics indicates amino acid(s) variation in epitope in comparison to the predicted epitope.

^b+++ indicates high binding affinity (<50 IC₅₀nm), ++ indicates medium binding affinity (>50 IC₅₀nm, <500 IC₅₀nm), + indicates poor binding affinity (>500 IC₅₀nm).

Table 5 Binding affinity scores of predicted epitopes and their variants to common HLA class II allele types prevalent in South Africa

Position	Predicted epitopes	Genotype of epitope	HLA II- DRB1 allele types ^a			
			HLA-DRB1*03:01	HLA-DRB1*04:01	HLA-DRB1*11:01	HLA-DRB1*15:01
E2						
507	YCFTPSPW	1a, 1b, 2, 3, 4, 5a, 6	+	+++	+	+++
695	NIVDTQYLY	5a	+	+++	+	+++
NS2						
938	LLHLGRLTG	5a	+	++	+++	+
1025	LLAPITAYA	1a, 3, 4, 5a,	+	++	+++	++
NS3						
1129	DLYLVTRHA	1a, 1b, 5a	+	+	+++	+
1131	YLVTRHADV	1a, 1b, 5a	+	+	+++	+
1391	LIFCHSKKK	1b, 2, 3, 4, 5a, 6	+	+	+++	+
1417	YYRGLDVAV	5a	+	+++	++	+
1464	FSLDPTFTI	1a, 1b, 2, 5a	+	+++	+++	+
1535	TTVRLRAYL	3, 5a, 6	+	+	+	+++
1562	FTGLTNIDA	5a	+	+++	++	+
1666	WAALAAYC	5a	+	+	+	+++
NS4B						
1774	YLAGLSTLP	1a, 1b, 2, 3, 4, 5a, 6	+	+++	+	+
1919	LIAFASRGN	1a, 1b, 2, 3, 4, 5a, 6	++	+++	+++	+++
1920	IAFASRGNH	1a, 1b, 2, 3, 4, 5a, 6	+	+	+++	+
NS5A						
1994	WLQAKLLPQ	5a	+	++	+++	+
2099	YHYITGVTQ	5a	+	++	+++	++
NS5B						
2450	LLRHHNLVY	1a, 3, 5a	+	++	++	+++
2579	LIVYDGLGV	2, 3, 4, 5a, 6	+	+	+	+++

^a+++ indicates high binding affinity (<50 IC₅₀nm), ++ indicates medium binding affinity (>50 IC₅₀nm, <500 IC₅₀nm), + indicates poor binding affinity (>500 IC₅₀nm).

immune responses [47]. A polyepitope-based strategy with multiple components combining core, E1, and E2 proteins; and conserved T-cell epitopes in the NS proteins has been suggested to be a good vaccine candidate for HCV [48].

High number of epitopes was predicted for HLA class II as compared to class I. The findings of this study are consistent with a study by Shehzadi et al. that predicted epitopes in genotype 3 from Pakistan. The study showed that majority of predicted epitopes were found in the NS3 protein for both HLA class I and HLA class II alleles and most of the epitopes were conserved among different genotypes [49]. Although the NS3 region is one of the conserved regions in HCV, variability in the nucleotide and amino acids has been reported by several studies in the same genotype and also in different genotypes [50,51]. A recent study that analysed 1568 NS3-protease sequences from genotypes 1–6 reported that the protease

amino acids sequence was moderately conserved and majority of the amino acids clustered in small regions. Of the 181 amino acids analysed 47% showed <1% variability among all HCV genotypes, and 17.1% amino acid positions showing >25.1% variability [51]. The NS3 is considered to be a good cellular target candidate for a therapeutic vaccine [52] since majority of the HCV viral epitopes recognized by CD8+ and CD4+ T-cells are located in the NS3 region [53-56]. The NS3 specific CD4+ and CD8+ T-cell responses were reported in patient responders to interferon therapy [57] and in spontaneous clearance of HCV [58].

Most of the predicted epitopes in the study sequence were found to be conserved across different HCV genotypes with a higher number of epitopes conserved at the anchor residues. The anchor residues are important for epitope high binding affinity to HLA [59]. Conserved epitopes might influence the immunogenic potential

since mutations within the epitopes can increase the chance of immune escape [60]. For a vaccine to be effective globally the selected epitopes must cover HLAs of different populations and it must also be conserved among different genotypes. The high mutation rates of viral epitopes and HLA polymorphisms are some of the challenges that are associated with the development of peptide vaccines [61]. Successful epitope vaccine design requires a broad knowledge of HCV genotype diversity. This will help in the proper selection of conserved HCV-specific T-cell epitopes that will help in avoiding HCV immune evasion [62]. This study attempted to ensure maximal coverage of HLA polymorphism and different genotypes by analyzing conserved epitopes considering different HLA alleles.

Majority of the epitopes predicted from HCV proteins isolated from South African genotype 5a were good binders against HLA alleles that are found worldwide. HLA is both polygenic and polymorphic, and the pool of HLA molecules differs for every individual. Different HLA alleles bind peptides with a particular sequence pattern [63]. For an HLA allele to be covered by a set of epitopes, at least one of the epitopes should be capable of inducing an immune response when bound to the corresponding HLA molecule [46]. The epitopes predicted in this study bind to many HLA alleles including the ones common in South Africa and can be used for designing good vaccine candidates that will eventually work in genetically diverse populations. *In-vitro* and *in-silico* studies have showed that HLA alleles preferentially bind to conserved regions of viral proteins in human viruses [64].

Very few epitopes were found to be experimentally true positive, however this can be due to the fact that most of the previous studies focused on genotype 1. A limitation of the study was a lack of *in-vivo* and *in-vitro* studies to confirm the predicted immunogenic epitopes, which will be the focus of future studies. However *in-silico* studies still provide the basis for designing good vaccine candidates.

In conclusion, the results of this study demonstrated antigenic T-cell epitopes that are conserved among genotypes and good HLA binders derived from genotype 5a sequences that can be good candidates for vaccine development. Predicted epitopes analysed in this study will contribute to the future design of an efficient vaccine with the use of conserved epitopes to avoid variation in genotypes and as such, it will be able to induce broad HCV specific immune responses. Conserved epitopes among different genotypes will be experimentally tested in the future to determine their involvement in immune response.

Methods

Ethical statement

The study was approved by the Medunsa Research and Ethics Committee of the Faculty of Health Sciences at the University of Limpopo as project no MREC/p/142/

2009:PG. The MREC is registered as an Independent Review Board with a reference no (IRB00005122).

Prediction of T-cell epitopes

Genotype 5a full-length sequences available in the GenBank and 6 of the near full length sequences generated from a previous study conducted by our group [24] were aligned and consensus sequences created using BioEdit [65] for the prediction of T-cell epitopes. For HLA class I, prediction for binding alleles was performed using ProPred I (<http://www.imtech.res.in/raghava/propred1/>) at a 4% default threshold by keeping the proteosome and immunoproteosome filters on at 5% threshold. ProPred 1 predicts antigenic epitopes for 47 HLA class I alleles [44]. For HLA class II, prediction was performed using ProPred (<http://www.imtech.res.in/raghava/propred/>) at a 3% default threshold. ProPred predicts antigenic epitopes for 51 HLA class II alleles [45].

Antigenicity of the epitopes

The Antigenicity score of all the predicted epitopes were analysed using Vaxijen v2.0 online antigen prediction (www.ddg-pharmfac.net/vaxijen/). Epitopes having antigenic score >0.5 were selected as antigenic. Vaxijen server performed well with 87% accuracy at a threshold of 0.5 antigenic score for viruses. Vaxijen v2.0 allows antigen classification based on the physicochemical properties of proteins without recourse to sequence alignment.

Epitope conservation analysis

All predicted epitopes were analyzed for conservation using the IEDB database (http://tools.immuneepitope.org/tools/conservancy/iedb_input) at a threshold of 100% conservation in comparison with 406, 221, 98, 33, 45, 45 randomly selected sequences from each of the HCV genotypes 1a, 1b, 2, 3, 4 and 6 respectively. The epitopes were considered conserved in another genotype if it shows 100% identity across the epitope in at least 70% of sequences in that genotype in the randomly selected sequences used in this study, downloaded from the public database. In addition, epitope variants that were conserved in at least 70% of the sequences were analysed for conservancy for anchor residues at positions 2 and 9 for HLA class I and positions 1, 4, 6 and 9 for HLA class II.

Validation of predicted epitopes

All the predicted epitopes were submitted to IEDB database (<http://www.immuneepitope.org/>) to confirm if they had been tested previously by other studies. The immuneepitope database contains experimentally confirmed data about antibody, T-cell epitopes, HLA binding, HLA restriction and HLA class.

Common South African HLA alleles

For epitope prediction binding to common HLA alleles found in South Africa, the IEDB epitope analysis tool (http://tools.immuneepitope.org/tools/conservancy/iedb_input) was used for HLA class I using the artificial neural network (ANN) algorithm [66] on the IEDB server, while for Class II ProPred (<http://www.imtech.res.in/raghava/propred/>) at a 3% default threshold was used. ProPred predicts antigenic epitopes for 51 HLA class II alleles [54]. The most common South African alleles were found in published literature [67].

Abbreviations

HCV: Hepatitis C virus; HLA: human leukocyte antigen.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MPG designed the study, performed the immunoinformatics analysis and drafted the manuscript. SGS and MJM guided the project and critically reviewed the manuscript. All the authors have read and approved the final manuscript.

Acknowledgements

This work was supported by grants from Medical Research Council, National Research Foundation, National Health Laboratory Service Research Trust, and The Stella & Paul Loewestein Charitable & Educational Trust in South Africa.

Received: 17 March 2014 Accepted: 14 October 2014

Published: 8 November 2014

References

- World Health Organization [WHO]: Guidelines for the screening care and treatment of persons with hepatitis C infection. 2014, 1–122 [<http://www.who.int/hiv/pub/hepatitis/hepatitis-c-guidelines/en/>]
- Pawlotsky JM: The nature of interferon-alpha resistance in hepatitis C virus infection. *Curr Opin Infect Dis* 2003, **16**:587–592.
- Ip PP, Nijman HW, Wilschut J, Daemen T: Therapeutic vaccination against chronic hepatitis C virus infection. *Antiviral Res* 2012, **96**:36–50.
- Ashfaq UA, Javed T, Rehman S, Nawaz Z, Riazuddin S: An overview of HCV molecular biology, replication and immune responses. *Virology* 2011, **8**:161.
- Singh R, Kaul R, Kaul A, Khan K: A comparative review of HLA associations with hepatitis B and C viral infections across global populations. *World J Gastroenterol* 2007, **13**:1770–1787.
- Thio CL, Nolt KR, Astemborski J, Vlahov D, Nelson KE, Thomas DL: Screening for hepatitis C virus in human immunodeficiency virus-infected individuals. *J Clin Microbiol* 2000, **38**(2):575–577.
- Fanning LJ, Kenny-Walsh E, Shanahan F: Persistence of hepatitis C virus in a white population: associations with human leukocyte antigen class 1. *Hum Immunol* 2004, **65**:745–751.
- McKiernan SM, Hagan R, Curry M, McDonald GS, Kelly A, Nolan N, Walsh A, Hegarty J, Lawlor E, Kelleher D: Distinct MHC class I and II alleles are associated with hepatitis C viral clearance, originating from a single source. *Hepatology* 2004, **40**:108–114.
- Neumann-Haefelin C, McKiernan S, Ward S, Viazov S, Spangenberg HC, Killinger T, Baumert TF, Nazarova N, Sheridan I, Pybus O, von Weizsäcker F, Roggendorf M, Kelleher D, Klennerman P, Blum HE, Thimme R: Dominant influence of an HLA-B27 restricted CD8+ T cell response in mediating HCV clearance and evolution. *Hepatology* 2006, **43**:563–572.
- Hong X, Yu RB, Sun NX, Wang B, Xu YC, Wu GL: Human leukocyte antigen class II DQB1*0301, DRB1*1101 alleles and spontaneous clearance of hepatitis C virus infection: a meta-analysis. *World J Gastroenterol* 2005, **11**:7302–7307.
- Lechner F, Wong DK, Dunbar PR, Chapman R, Chung RT, Dohrenwend P, Robbins G, Phillips R, Klennerman P, Walker BD: Analysis of successful immune responses in persons infected with hepatitis C virus. *J Exp Med* 2000, **191**:1499–1512.
- Spada E, Mele A, Berton A, Ruggeri L, Ferrigno L, Garbuglia AR, Perrone MP, Girelli G, Del Porto P, Piccolella E, Mondelli MU, Amoroso P, Cortese R, Nicosia A, Vitelli A, Folgori A: Multispecific T cell response and negative HCV RNA tests during acute HCV infection are early prognostic factors of spontaneous clearance. *Gut* 2004, **53**:1673–1681.
- Lavillette D, Morice Y, Germanidis G, Donot P, Soulier A, Pagkalos E, Sakellariou G, Intrator L, Bartosch B, Pawlotsky JM, Cosset FL: Human serum facilitates hepatitis C virus infection, and neutralizing responses inversely correlate with viral replication kinetics at the acute phase of hepatitis C virus infection. *J Virol* 2005, **79**(10):6023–6034.
- Petrovic D, Dempsey E, Doherty DG, Kelleher D, Long A: Hepatitis C virus-T-cell responses and viral escape mutations. *Eur J Immunol* 2012, **42**:17–26.
- Shoukry NH, Grakoui A, Houghton M, Chien DY, Ghayeb J, Reimann KA, Walker CM: Memory CD8+ T cells are required for protection from persistent hepatitis C virus infection. *J Exp Med* 2003, **197**:1645–1655.
- Klennerman P, Thimme R: T cell responses in hepatitis C: the good, the bad and the unconventional. *Gut* 2012, **61**:1226–1234.
- Brown RJ, Tarr AW, McClure CP, Juttla VS, Tagiuri N, Irving WL, Ball JK: Cross-genotype characterization of genetic diversity and molecular adaptation in hepatitis C virus envelope glycoprotein genes. *J Gen Virol* 2007, **88**:458–469.
- Patronov A, Doytchinova I: T-cell epitope vaccine design by immunoinformatics. *Open Biol* 2013, **3**:120139.
- Dimitrov I, Flower DR, Doytchinova I: Improving in Silico Prediction of Epitope Vaccine Candidates by Union and Intersection of Single Predictors. *W J V* 2011, **1**:15–22.
- Gedezha MP, Selabe SG, Kyaw T, Rakgole NJ, Blackard JT, Mphahlele MJ: Introduction of new subtypes and variants of hepatitis C virus genotype 4 in South Africa. *J Med Virol* 2012, **84**:601–607.
- Antaki N, Craxi A, Kamal S, Mouchari R, Van der Merwe S, Haffar S, Gadano A, Zein N, Lai CL, Pawlotsky JM, Heathcote EJ, Dusheiko G, Marcellin P: The neglected hepatitis C virus genotypes 4, 5 and 6: an international consensus report. *Liver Int* 2010, **30**:342–355.
- Bukh J, Apparg CL, Engle R, Govindarajan S, Hegerich PA, Tellier R, Wong DC, Elkins R, Kew MC: Experimental infection of chimpanzees with hepatitis C virus of genotype 5a: genetic analysis of the virus and generation of a standardized challenge pool. *J Infect Dis* 1998, **178**:1193–1197.
- Chamberlain RW, Adams NJ, Taylor LA, Simmonds P, Elliott RM: The complete coding sequence of hepatitis C virus genotype 5a, the predominant genotype in South Africa. *Biochem Biophys Res Commun* 1997, **236**:44–49.
- Gedezha MP, Selabe SG, Blackard JT, Kyaw T, Mphahlele MJ: Near full-length genome analysis of HCV genotype 5 strains from South Africa. *Infect Genet Evol* 2014, **21**:118–123.
- Himoudi N, Abraham J, Fournillier A, Lone YC, Joubert A, Op De Beeck A, Freida D, Lemonnier F, Kieny MP, Inchauspé G: Comparative vaccine studies in HLA-A2.1-transgenic mice reveal a clustered organization of epitopes presented in hepatitis C virus natural infection. *J Virol* 2002, **76**:12735–12746.
- Ohno S, Moriya O, Yoshimoto T, Hayashi H, Akatsuka T, Matsui M: Immunogenic variation between multiple HLA-A*0201-restricted, Hepatitis C Virus-derived epitopes for cytotoxic T lymphocytes. *Viral Immunol* 2006, **19**:458–467.
- Engler OB, Schwendener RA, Dai WJ, Wölk B, Pichler W, Moradpour D, Brunner T, Cerny A: A liposomal peptide vaccine inducing CD8+ T cells in HLA-A2.1 transgenic mice, which recognise human cells encoding hepatitis C virus (HCV) proteins. *Vaccine* 2004, **23**:58–68.
- Ibe M, Sakaguchi T, Tanaka K, Saito S, Yokota S, Tanaka T, Shimotohno K, Chujoh Y, Shiratori Y, Omata M, Miwa K, Takiguchi M: Identification and characterization of a cytotoxic T cell epitope of hepatitis C virus presented by HLA-B*3501 in acute hepatitis. *J Gen Virol* 1998, **79**:1735–1744.
- Giugliano S, Oezkan F, Bedrejowski M, Kudla M, Reiser M, Viazov S, Scherbaum N, Roggendorf M, Timm J: Degree of cross-genotype reactivity of hepatitis C virus-specific CD8+ T cells directed against NS3. *Hepatology* 2009, **50**:707–716.
- Park S, Shin E, Capone S, Caggiari L, Re VD, Nicosia A, Folgori A, Rehmann B: Successful vaccination induces multifunctional memory T-cell precursors associated with early control of hepatitis C virus. *Gastroenterology* 2012, **143**:1048–1060.

31. Boucherma R, Kridane-Miledi H, Bouziat R, Rasmussen M, Gatar T, Langa-Vives F, Lemerrier B, Lim A, Bérard M, Benmohamed L, Buus S, Rooke R, Lemonnier FA: **HLA-A*01:03, HLA-A*24:02, HLA-B*08:01, HLA-B*27:05, HLA-B*35:01, HLA-B*44:02, and HLA-C*07:01 monochain transgenic/H-2 class I null mice: novel versatile preclinical models of human T cell responses.** *J Immunol* 2013, **191**:583–593.
32. Doi H, Hiroishi K, Shimazaki T, Eguchi J, Baba T, Ito T, Matsumura T, Nozawa H, Morikawa K, Ishii S, Hiraide A, Sakaki M, Imawari M: **Magnitude of CD8 T-cell responses against hepatitis C virus and severity of hepatitis do not necessarily determine outcomes in acute hepatitis C virus infection.** *Hepatol Res* 2009, **39**:256–265.
33. Wentworth PA, Sette A, Celis E, Sidney J, Southwood S, Crimi C, Stitely S, Keogh E, Wong NC, Livingston B, Alazard D, Vitiello A, Grey HM, Chisari FV, Chesnut RW, Fikes J: **Identification of A2-restricted hepatitis C virus-specific cytotoxic T lymphocyte epitopes from conserved regions of the viral genome.** *Int Immunol* 1996, **8**:651–659.
34. Wang S, Buchli R, Schiller J, Gao J, VanGundy RS, Hildebrand WH, Eckels DD: **Natural epitope variants of the hepatitis C virus impair cytotoxic T lymphocyte activity.** *World J Gastroenterol* 2010, **16**:1953–1969.
35. Fitzmaurice K, Petrovic D, Ramamurthy N, Simmons R, Merani S, Gaudieri S, Sims S, Dempsey E, Freitas E, Lea S, McKiernan S, Norris S, Long A, Kelleher D, Klenerman P: **Molecular footprints reveal the impact of the protective HLA-A*03 allele in hepatitis C virus infection.** *Gut* 2011, **60**:1563–1571.
36. Scognamiglio P, Accapezzato D, Casciaro MA, Cacciani A, Artini M, Bruno G, Chiric ML, Sidney J, Southwood S, Abrignani S, Sette A, Barnaba V: **Presence of effector CD8+ T cells in hepatitis C virus-exposed healthy seronegative donors.** *J Immunol* 1999, **162**:6681–6689.
37. Chang KM, Gruener NH, Southwood S, Sidney J, Pape GR, Chisari FV, Sette A: **Identification of HLA-A3 and -B7-restricted CTL response to hepatitis C virus in patients with acute and chronic hepatitis C.** *J Immunol* 1999, **162**:1156–1164.
38. Sreenarasimhaiah J, Jaramillo A, Crippin J, Lisker-Melman M, Chapman WC, Mohanakumar T: **Concomitant augmentation of type 1 CD4+ and CD8+ T-cell responses during successful interferon-alpha and ribavirin treatment for chronic hepatitis C virus infection.** *Hum Immunol* 2003, **64**:497–504.
39. Matsui M, Moriya O, Belladonna ML, Kamiya S, Lemonnier FA, Yoshimoto T, Akatsuka T: **Adjuvant activities of novel cytokines, interleukin-23 (IL-23) and IL-27, for induction of hepatitis C virus-specific cytotoxic T lymphocytes in HLA-A*0201 transgenic mice.** *J Virol* 2004, **78**:9093–9104.
40. Mizukoshi E, Nascimbeni M, Blaustein JB, Mihalik K, Rice CM, Liang TJ, Feinstone SM, Rehermann B: **Molecular and immunological significance of chimpanzee major histocompatibility complex haplotypes for hepatitis C virus immune response and vaccination studies.** *J Virol* 2002, **76**:6093–6103.
41. Zhang J, Yamada O, Yoshida H, Iwai T, Araki H: **Autogenous translational inhibition of core protein: implication for switch from translation to RNA replication in hepatitis C virus.** *Virology* 2002, **293**:141–150.
42. Matsueda S, Yamada A, Takao Y, Tamura M, Komatsu N, Yutani S, Ide T, Sata M, Itoh K: **A new epitope peptide derived from hepatitis C virus 1b possessing the capacity to induce cytotoxic T-lymphocytes in HCV1b-infected patients with HLA-A11, -A31, and -A33.** *Cancer Immunol Immunother* 2007, **56**:1359–1366.
43. De Groot AS, Sbai H, Aubin CS, McMurry J, Martin W: **Immuno-informatics: Mining genomes for vaccine components.** *Immunol Cell Biol* 2002, **80**:255–269.
44. Singh H, Raghava GP: **ProPred1: prediction of promiscuous MHC Class-I binding sites.** *Bioinformatics* 2003, **19**:1009–1014.
45. Singh H, Raghava GP: **ProPred: prediction of HLA-DR binding sites.** *Bioinformatics* 2001, **17**:1236–1237.
46. Toussaint NC, Dönnes P, Kohlbacher O: **A mathematical framework for the selection of an optimal set of peptides for epitope-based vaccines.** *PLoS Comput Biol* 2008, **4**:e1000246.
47. Zeng R, Li G, Ling S, Zhang H, Yao Z, Xiu B, He F, Huang R, Wei L: **A novel combined vaccine candidate containing epitopes of HCV NS3, core and E1 proteins induces multi-specific immune responses in BALB/c mice.** *Antiviral Res* 2009, **84**:23–30.
48. Hu CT: **Vaccine Development for Hepatitis C: Lessons from the Past Turn into Promise for the Future.** *Tzu Chi Med J* 2005, **17**:61–74.
49. Shehzadi A, Ur Rehman S, Idrees M: **Promiscuous prediction and conservancy analysis of CTL binding epitopes of HCV 3a viral proteome from Punjab Pakistan: an in silico approach.** *Viral J* 2011, **8**:55.
50. Vallet S, Gouriou S, Nousbaum JB, Legrand-Quillien MC, Goudeau A, Picard B: **Genetic heterogeneity of the NS3 protease gene in hepatitis C virus genotype 1 from untreated infected patients.** *J Med Virol* 2005, **75**(4):528–537.
51. Cento V, Mirabelli C, Salpini R, Dimonte S, Artese A, Costa G, Mercurio F, Svicher V, Parrotta L, Bertoli A, Ciotti M, Di Paolo D, Sarrecchia C, Andreoni M, Alcaro S, Angelico M, Perno CF, Ceccherini-Silberstein F: **HCV genotypes are differently prone to the development of resistance to linear and macrocyclic protease inhibitors.** *PLoS One* 2012, **7**(7):e39652.
52. Torresi J, Johnson D, Wedemeyer H: **Progress in the development of preventive and therapeutic vaccines for hepatitis C virus.** *J Hepatol* 2011, **54**:1273–1285.
53. Eckels DD, Bian T, Gill JC, Sønderstrup G: **Epitopes of the NS3 protein of hepatitis C virus: recognition in HLA-DR4 transgenic mice.** *Immunol Cell Biol* 2002, **80**:106–112.
54. Day CL, Lauer GM, Robbins GK, McGovern B, Wurcel AG, Gandhi RT, Chung RT, Walker BD: **Broad specificity of virus-specific CD4+ T-helper-cell responses in resolved hepatitis C virus infection.** *J Virol* 2002, **76**:12584–12595.
55. Hakamada T, Funatsuki K, Morita H, Ugajin T, Nakamura I, Ishiko H, Matsuzaki Y, Tanaka N, Imawari M: **Identification of novel hepatitis C virus-specific cytotoxic T lymphocyte epitopes by ELISpot assay using peptides with human leukocyte antigen-A*2402-binding motifs.** *J Gen Virol* 2004, **85**:1521–1531.
56. Wertheimer AM, Miner C, Lewinsohn DM, Sasaki AW, Kaufman E, Rosen HR: **Novel CD4+ and CD8+ T-cell determinants within the NS3 protein in subjects with spontaneously resolved HCV infection.** *Hepatology* 2003, **37**:577–589.
57. Vertuani S, Bazzaro M, Gualandi G, Micheletti F, Marastoni M, Fortini C, Canella A, Marino M, Tomatis R, Traniello S, Gavioli R: **Effect of interferon-alpha therapy on epitope-specific cytotoxic T lymphocyte responses in hepatitis C virus-infected individuals.** *Eur J Immunol* 2002, **32**:144–154.
58. Smyk-Pearson S, Tester IA, Lezotte D, Sasaki AW, Lewinsohn DM, Rosen HR: **Differential antigenic hierarchy associated with spontaneous recovery from hepatitis C virus infection: implications for vaccine design.** *J Infect Dis* 2006, **194**:454–463.
59. Hammer J, Belunis C, Bolin D, Papadopoulos J, Walsky R, Higelin J, Danho W, Sinigaglia F, Nagy ZA: **High-affinity binding of short peptides to major histocompatibility complex class II molecules by anchor combinations.** *Proc Natl Acad Sci U S A* 1994, **91**:4456–4460.
60. Yusim K, Kesmir C, Gaschen B, Addo MM, Altfeld M, Brunak S, Chigaev A, Detours V, Korber BT: **Clustering patterns of cytotoxic T-lymphocyte epitopes in human immunodeficiency virus type 1 (HIV-1) proteins reveal imprints of immune evasion on HIV-1 global variation.** *J Virol* 2002, **76**:8757–8768.
61. Vider-Shalit T, Raffaelli S, Louzoun Y: **Virus-epitope vaccine design: Informatic matching the HLA-I polymorphism to the virus genome.** *Mol Immunol* 2007, **44**:1253–1261.
62. Molero-Abraham M, Lafuente EM, Flower DR, Reche PA: **Selection of conserved epitopes from hepatitis C virus for pan-population stimulation of T-cell responses.** *Clin Dev Immunol* 2013, **2013**:601943.
63. Doytchinova IA, Guan P, Flower DR: **Epiln: a server for multistep T cell epitope prediction.** *BMC Bioinformatics* 2006, **7**:131.
64. Hertz T, Nolan D, James I, John M, Gaudieri S, Phillips E, Huang JC, Riadi G, Mallal S, Jojic N: **Mapping the landscape of host-pathogen coevolution: HLA class I binding and its relationship with evolutionary conservation in human and viral proteins.** *J Virol* 2011, **85**(3):1310–1321.
65. Hall TA: **BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT.** *Nucl Acids Symp Ser* 1999, **41**:95–98.
66. Nielsen M, Lundegaard C, Worning P, Laue-moller SL, Lambert K, Buus S, Brunak S, Lund O: **Reliable prediction of T-cell epitopes using neural networks with novel sequence representations.** *Protein Sci* 2003, **12**:1007–1017.
67. Paximadis M, Mathebula TY, Gentle NL, Vardas E, Colvin M, Gray CM, Tiemessen CT, Puren A: **Human leukocyte antigen class I (A, B, C) and II (DRB1) diversity in the black and caucasian South African population.** *Hum Immunol* 2012, **73**:80–92.

doi:10.1186/1743-422X-11-187

Cite this article as: Gedezha *et al.*: Prediction of T-cell epitopes of hepatitis C virus genotype 5a. *Virology Journal* 2014 **11**:187.