

RESEARCH

Open Access

Gene expression profiling of CD4⁺ T cells in treatment-naive HIV, HCV mono- or co-infected Chinese

Lina Yi^{1,3†}, Jin Zhao^{1,2†}, Jing Lu^{3†}, Ying Chen¹, Lin Chen², Jinqian Cheng², Yan Sun⁴, Zhi Li⁴, Ruotin Men⁵, Li Yang⁵, Hsiangfu Kung¹, Zhengrong Yang^{2*} and Ming-liang He^{1,6*}

Abstract

Background: Because of the shared transmission routes, co-infection with human immunodeficiency virus (HIV) and hepatitis C virus (HCV) is very common. Accumulated clinical evidence showed that one could alter the infectious course of the other virus in HIV and HCV co-infected individuals. However, little is known on the molecular basis of HIV/HCV interactions and their modulations on hosts.

Methods: In this study, treatment-naive HIV, HCV mono-/co-infected individuals with CD4⁺ T cell counts >300/ μ l were recruited and their gene expression profiles were investigated by microarray assays. The differentially expressed genes were identified and validated by quantitative real-time PCR (qRT-PCR). To further understand the biological meanings of the gene expression profiles in these three groups, GSEA analysis (version 2.0, Broad Institute <http://www.broad.mit.edu/gsea>) was performed.

Results: By gene set enrichment analysis, we revealed that gene sets of cell cycle progression, innate immune response and some transcription factors in CD4⁺ T cells were mainly affected by HIV; while genes associated with GPCR signaling were the major targets of HCV. Metabolic pathways were modulated by both HCV and HIV viruses.

Conclusions: This study for the first time offers gene profiling basis for HCV/HIV mono-/co-infections in human beings. HIV infection displayed the great impact on transcription profile of CD4⁺ T cells in HIV/HCV co-infected individuals. Genes related to cell cycle arrest were significantly mediated by HIV which may lead to dysfunction of CD4⁺ T cells and acceleration of HCV-related disease progression in the co-infections.

Keywords: HIV, HCV, Co-infection, Microarray, CD4⁺ T cells

Background

Co-infection with human immunodeficiency virus (HIV) and hepatitis C virus (HCV) is very common, presumably due to their shared transmission routes (e.g., drug injection, sex behaviour) [1,2]. It is estimated that over 5 million people are HCV/HIV co-infected worldwide [2]. The prevalence of HCV/HIV co-infection varies widely in different risk groups, with an especially high rate among HIV-positive intravenous drug users (IDUs) [3].

In the United States and China, 72% to 90% IDUs in HIV-infected persons were co-infected with HCV [4].

Accumulated evidence showed that one could alter the course of infection of the other virus in HIV and HCV co-infected individuals. On the one hand, a higher rate of HCV viral persistence and increased viral load are more common in the HCV/HIV co-infected patients than in HCV mono-infected ones [5]. Moreover, it was suggested that HIV could accelerate the progression of HCV-associated liver diseases, including fibrosis, cirrhosis and end-stage liver disease [6-8]. On the other hand, an acute HCV infection normally raises the HIV-induced viremia in persons with otherwise well-controlled illness, and accelerates the progression to AIDS and death by impairing immune reconstitution [9]. Besides, the risk of

* Correspondence: zryang2012@gmail.com; mlhe7788@gmail.com

[†]Equal contributors

²Shenzhen Center for Disease Control and Prevention, Shenzhen, China

¹Stanley Ho Center for Emerging Infectious Diseases, and Li Ka Shing Institute of Health Sciences, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China

Full list of author information is available at the end of the article

hepatotoxicity from highly active antiretroviral treatment (HAART) may be increased in the co-infected patients as compared to the HIV mono-infected patients, thus lead to a decreased tolerability for the anti-HIV treatment [10].

Mechanisms of the accelerated progression of diseases in the co-infected patients are not yet well understood. Analysis of gene expression profiles in the HCV/HIV mono-infected or co-infected patients would provide a unique opportunity to understand the mechanisms. It is reported that factors, such as direct viral effects, the immunologic alterations especially the specific T-cell response, have played important roles in the disease progression [11-13]. CD4⁺ T cells, the major HIV target cells, also play an essential role in HCV clearance [14,15]. In this study, we performed the microarray studies to analysis the gene expression profiles of CD4⁺ T cells from treatment-naïve HIV/HCV mono- and co-infected individuals. Moreover, via gene set enrichment analysis (GSEA), a network of enriched pathways related to the pathogenesis of disease progression in the co-infected patients was identified. To our knowledge, this is the first study to analyze the impact of HIV/HCV co-infection at gene transcription level, and our data may offer new insight into understanding the interplay during HIV and HCV co-infection.

Results

Differentially expressed genes

Microarray analysis was performed to identify the altered transcripts in 3 study groups with HIV, HCV mono-infection and HCV/HIV co-infections, and the raw data has been deposited in the ArrayExpress database (access number: E-MEXP-3601). Pairwise comparisons from the three study groups (HCV-mono versus HIV-mono, HCV/HIV co-infected versus HCV-mono, HCV/HIV co-infected versus HIV-mono) were carried out. Differentially expressed (DE) transcript identifiers (DETI) with >2 fold change and $p < 0.05$ were identified for each comparison. The number of DETIs identified in each comparison is listed in Table 1. The detailed information of each DETI in each comparison was listed in Additional file 1: Table S2.

To identify the important categories from the DE genes, functions of each DE gene were inspected. Of 54

DE genes identified in comparison between HCV and HIV mono-infected groups, 24 showed significant matches with the known genes. Functional analysis revealed that genes involved in immune system development, immune response (CXCL10, OAS3, CD38, SERPING1, CCR1, FCGR3A and IFI44L) and G-protein coupled receptor (GPCR) signaling pathway (P2RY13, GPR56, CX3CR1 and FFAR2) were differently affected by these two viruses. In the case of HCV/HIV co-infection versus HCV mono-infection, 72 genes with known function were identified. Many of them (16/72) played important roles in stimulus response and immune system process, including: RSAD2, IL6, C1QC, FCGR1C, LILRA3, IGSF6, IL1RN, OAS3. Besides, genes involved in regulating locomotory behavior such as CCRL2, CX3CR1, FPR2 and IL8RA were also shown different expression profiles between these two groups. In comparison between the HCV/HIV co-infections and the HIV mono-infections, only 7 were identified to be known genes. 3 were small nucleolar RNAs, and the remaining 4 were TNFAIP6, AQP9, IL6 and PTX3. The later four genes play important role in immune response and involve in many human diseases, such as cancers and diseases caused by a variety of virus infections.

Gene set enrichment analysis


To unravel the biological mechanisms differentiating between the HCV/HIV mono- and the co-infected groups, pairwise comparisons using GSEA were performed for all the three groups (HCV-mono versus HIV-mono, HCV/HIV co-infected versus HCV-mono, HCV/HIV co-infected versus HIV-mono). Rather than single DE genes, GSEA evaluates microarray data at the biological pathway level by performing unbiased global searches for genes that are coordinately regulated in predefined gene sets. The number of significantly enriched gene sets ($FDR < 0.01$) in each pairwise comparison is listed in Table 1. And pathways identified in gene set enrichment analysis (GSEA, $FDR < 0.05$) were shown in Additional file 2: (Table S3).

Global gene expression profiles between HIV mono- and HCV/HIV co-infected groups

We first applied GSEA to identify functional gene sets distinct between the HIV mono- and the HCV/HIV

Table 1 Number of differentially expressed transcript identifiers (fold change > 2 and P < 0.05) and enriched gene sets (FDR < 0.01) in pair wise comparisons for CD4⁺ T cells

	HCV-mono		HCV/HIV co-infected		HCV/HIV co-infected	
	versus		versus		versus	
	HIV-mono		HCV-mono		HIV-mono	
	up	down	up	down	up	down
Differentially expressed transcript identifiers	19	35	109	79	44	7
Enriched gene sets	6	44	9	10	0	2



GPCR signaling pathway was up-regulated in HCV mono-infected individuals no matter compared with HIV or HCV/HIV-infected individuals

Besides the above mentioned gene sets, there still existed some other pathways that showed high overlap between these two pairwise comparisons (HCV-mono versus HIV-mono, HCV/HIV co-infected versus HCV-mono). Among them, the gene sets involved in signal transduction were top listed. Contrary to the cell cycle category, these gene sets were up-regulated in HCV mono-infected individuals. Further analysis revealed that nearly these entire gene sets could be directly or indirectly associated with GPCR signaling pathway.

Validation of differentially expressed genes

To confirm the DE genes from the microarray analysis, the mRNA levels from each paired comparison were selected and quantitated by qRT-PCR. These genes included P2RY13, Mx1, IL6, PTX3, GPR56, OAS1, CX3CR1, USP18 (Figure 4) and FCGR3A, CCR1, TLR4, IFI44L, CD38 (Supplemental Figure 1). The RNA isolated

from the CD4⁺ T cells of each individual (16 individuals in each group) was used. And the blood samples from un-infected males with matched ages were collected as control. Expression levels of these genes in four groups were all evaluated (Figure 4) and compared with the results obtained by microarray (Table 2). The qRT-PCR results showed that the expressing profiles of 6 genes (PTX3, IL6, P2RY13, OAS1, CX3CR1 and USP18) exactly matched with the observation in microarray assays; another 2 genes (Mx1 and GPR56) displayed partial similarity at least in one express pattern out of three pairwise comparisons. These results largely confirmed the data from the microarray assays. Besides, the mRNA levels of all the selected genes in un-infected persons with matched ages were also analyzed. As shown in Figure 4 and Table 2, the expression levels of CX3CR1, PTX3, MX1 and P2RY13 decreased significantly in infected groups as compared to the uninfected healthy group; while the expression level of OAS1 elevated in the infected groups. There is one gene (USP18) showing altered expressions only in HCV-infections. For IL6, lower expression levels




Figure 3 Enrichment plot and heat map for the gene set of immune system identified in comparison between HCV/HIV co-infected group and HCV mono-infected group. **A.** Enrichment plot for CD4⁺ T cells from comparison between HCV/HIV co-infected group and HCV mono-infected group. Bottom, plot of the ranked list of all genes. Y axis, value of the ranking metric; X axis, the rank for all genes. Genes whose expression levels are most closely associated with the HCV mono-infected group or HCV/HIV co-infected group get the highest metric scores with positive or negative sign, and are located at the left or right edge of the list. Middle, the location of genes from the gene set immune system within the ranked list. Top, the running enrichment score for the gene set as the analysis walks along the ranked list. The score at the peak of the plot is the enrichment score (ES) for this gene set and those genes appear before or at the peak are defined as core enrichment genes for this gene set. **B.** Heat map of the genes within the gene set of immune system corresponding to **A.** The genes that contribute most to the ES, i.e., genes that appear in the ranked list before or at the peak point of ES, are defined as core enrichment genes and highlighted by the red rectangle. Rows, genes; columns, samples. Range of colors (red to blue) shows the range of expression values (high to low).

were observed both in the HCV and HIV infections. Other immune related genes like FCGR3A and CCR1 were significantly upregulated while CD38 were decreased in co-infected group as compared to the mono infected or uninfected groups (Additional file 3: Figure 1).


Discussion

Co-infection with HIV and HCV is common, and increasing evidence indicated that each can alter the course of infection of the other one [16]. However, little is known on the molecular basis of HIV and HCV interactions and their modulations on host responses. In this study, the gene expression profiles of CD4⁺ T cells in the treatment-naive HCV, HIV mono-infected and the HCV/HIV co-infected individuals were evaluated. For each cell subset, pairwise comparisons were performed and differentially expressed genes and significantly altered pathways were identified.

By using the GSEA analysis, groups of comprehensive pathways were identified. At first glance, it was obvious

to notice that the major altered gene sets identified in comparison between the HCV/HIV co-infections and HCV mono-infections were nearly same with those identified in comparison between HIV and HCV mono-infections. Considering that CD4⁺ T cell was the major target of HIV and only a little specific CD4⁺ T cells was detected in the HCV infected individuals [16,17], it was reasonable that HCV only contribute a little to the global gene changes of CD4⁺ T cells in HCV/HIV co-infections (Table 1).

One significant proportion of differentially expressed genes was associated with cell cycle which was significant up-regulated in both the HIV mono- and HCV/HIV co-infected individuals as compared with HCV mono-infected individuals (Figure 2). Functional analysis revealed that most of these gene sets involved in G1/S and G2/M transitions. In relate to G1/S transition, till now only one study has reported this phenomenon and the reason why HIV prefers to drive cells out of G1 to S phase is not clear yet [18]. One possible explanation is



that the G1/S transition may regulate the latency of HIV [19]. The promotion of G1/S transition may facilitate HIV integration as indicated by the up-regulation of genes engaged mainly in DNA repair. The latter has been assumed to be required for the integration of HIV

[20,21]. Another most affected phase was the G2/M transition [22,23]. As shown in Figure 2, 16 out of 33 genes that associated with G2/M transition were up-regulated in the HIV infected individuals, and most of them play important roles in the maintenance of

Table 2 Quantitative real time PCR validations of differentially expressed genes

Gene symbol	Method	HIV + HCV vs HCV	HIV + HCV vs HIV	HCV vs HIV	HCV vs None-infection	HIV vs None-infection	HCV + HIV vs None-infection
IL6	MA	Up	Up	NS	ND	ND	ND
	qPCR	Up	Up	NS	Down	Down	NS
PTX3	MA	Up	Up	NS	ND	ND	ND
	qPCR	Up	Up	NS	Down	Down	Down
Mx1	MA	Up	NS	Down	ND	ND	ND
	qPCR	Down	Down	Down	Down	Down	Down
USP18	MA	Up	NS	Down	ND	ND	ND
	qPCR	Up	NS	Down	Down	NS	NS
P2RY13	MA	Up	NS	Down	ND	ND	ND
	qPCR	Up	NS	Down	Down	Down	Down
GPR56	MA	Up	NS	Down	ND	ND	ND
	qPCR	NS	NS	Down	NS	NS	NS
OAS1	MA	Up	NS	NS	ND	ND	ND
	qPCR	Up	NS	NS	Up	Up	Up
CX3CR1	MA	Up	NS	Down	ND	ND	ND
	qPCR	Up	NS	Down	Down	Down	Down

NOTE. VS, versus; NS, not significantly changed; ND, not detected; qPCR, quantitative real-time polymerase chain reaction; MA, microarray analysis.

centrosome normality and integrity. The Vpr (viral protein R) of HIV was reported to induce an accumulation of multiple centrosome-like structures in human cells that lead to cell cycle arrest or delay in the G2 phase [24], thus positively support viral replication or facilitate viral spread by triggering cell death [25]. For HIV/HCV co-infection, the cell cycle arrest in G2 phase induced by HIV would lead to a depletion of CD4⁺ T cells [25], and may partially result in a less efficient in HCV clearance [26].

The second group of significantly up-regulated gene sets in both HIV mono- and HCV/HIV co-infections was genes related to innate immune response as compared with HCV mono-infections (Figure 3). In chronic infection, such as HIV, continuous ongoing innate immune responses may contribute more to disease progression rather than to limit viral replication [26]. Although the mechanisms are very complicated, functional analysis revealed that a group of overlapped genes were engaged in NF- κ B regulation among these different innate immune pathways. This finding was also supported by other studies that NF- κ B was mediated by Vpr and played a major role in HIV gene expression and pro-inflammatory cytokines induction [27-30].

The third group of gene sets that is usually modulate by virus is the host cell transcription apparatus [31]. By providing direct gene pattern, our results confirmed that this strategy was also exploited both in the HIV mono- and HCV/HIV co-infected individuals. As seen in Figure 1C, many of the critical components that engaged in mRNA formation, elongation and maturation were up-regulated in the HIV mono- and HCV/HIV co-infected individuals. These implied that through up-regulating the mRNA editing genes, the HIV virus could facilitate the production of matured transcripts which were needed for its own translation.

Gene sets in metabolic pathways, including carbohydrate, lipid, fatty acid, amino acid, nucleotide and even vitamin metabolisms, were all significantly changed in the co-infected group. Both of these two viruses have shown abilities to modulate partial of these pathways to facilitate their infections [32-34]. The abnormalities in the whole metabolic net in co-infections may be a result of HIV-HCV interaction. By complement to each other, the damaging effects would be definitely aggravated, thus lead a more rapid disease progression in co-infections.

Results from our study also detected several pathways up-regulated in HCV infections. All of these pathways were directly or indirectly relate to pathway of GPCR signaling (Figure 1). By modulation several major effectors such as adenylyl cyclase, phospholipase C and the mitogen activated protein kinases (MAPKs), GPCR are involved in many diverse signaling events including visual sense, smell, immune system regulation and inflammation [35,36]. The significance of HCV on modulating GPCR signaling is not

well understood yet. We speculate that the activation of GPCR signaling as exemplified by upregulation of chemokines (e.g., CCL22) may play important roles in lymphocytes chemotaxis, which could promote the activated lymphocytes to inflammatory sites and facilitate viral clearance [37]. However, an inappropriate activation of chemotaxis may also lead to unexpected damage of un-infected cells and accelerate the disease progression. Activation of GPCR pathways would lead to an increase in amount and duration of intracellular cyclic adenosine 3', 5'-monophosphate (cAMP) levels [38]. As an important regulator of immune cells, cAMP has a dual and opposite role during HIV infection [39]. cAMP can limit viral entry and replication [40,41], while it may also reduce HIV-specific antiviral immune responses and promote T cell dysfunctions [42,43]. The exact role of HCV-activated GPCR signaling on CD4⁺ T cells function and HIV infection needed to be further determined.

Conclusion

This study for the first time analyzed the impact of HIV/HCV co-infection at gene transcription level. By comparing with HCV and HCV/HIV co-infection, HIV infection displayed the great impact on gene expression profile of CD4⁺ T cells. Genes related to cell cycle arrest and T cell dysfunction were significantly mediated in HIV and HIV/HCV infected individuals which may practically explain the acceleration of HCV related disease progression by HIV infection and control of HIV replication might improve HCV specific CD4⁺ T-cell response in the HCV/HIV co-infected individuals.

Methods

Treatment-naive HCV/HIV mono-/co-infected individuals

A male population of Chinese was recruited from an ongoing voluntary-based HIV/AIDS surveillance study in Shenzhen, from September 2009 to December 2010. Written content was obtained from the participants. Ethics approval was obtained from The Joint Chinese University of Hong Kong–New Territories East Cluster Clinical Research Ethics Committee. All participants have been screened for HIV (Beijing Wantai Biological Pharmacy Enterprise CO., LTD, Beijing), HCV (Beijing Wantai Biological Pharmacy Enterprise CO., LTD, Beijing) and HBV (Beijing Wantai Biological Pharmacy Enterprise CO., LTD, Beijing) antigens and antibodies by standard ELISA analysis recommended by The Center of Disease Control and Prevention of China (guidelines of CDC, China). HIV infection was further confirmed by western blot (HIV Blot 2.2 WB; Genelabs Diagnostics, Singapore) following the National Guideline for Detection of HIV/AIDS of China. In order to rule out the confounding factors including treatment responses and drug resistant bias, all HIV + samples were tested for CD4 counts with flow

cytometry, and drug resistant mutations by genotyping analysis. All enrolled HIV+ individuals are antiretroviral-naïve with CD4 counts ≥ 300 cells/ μ l and without baseline drug resistant mutation. Based on the stringent criteria described above, 24 samples from each infected group (HIV mono-infected, HCV mono-infected and HIV/HCV co-infected group) were selected in this study. For microarray analysis, each group contained 3 biological replicates with 8 individuals in each replicates. To validate our findings by quantitative real-time PCR (qRT-PCR), 24 healthy individuals were further enrolled. Their demographic and epidemiological data have been documented in Additional file 4: Table S1. The average age was similar in all the study groups (29.5, 33.0, 31.4 and 27.5 years for HIV mono-infection group, HCV mono-infection group, HIV/HCV co-infection group and non-infected group, respectively). The CD4 counts were 433 ± 101 , 839 ± 234 , 524 ± 219 and 651 ± 210 (mean \pm SD, cells/ μ l) for HIV infected, HCV infected, HIV/HCV co-infected and un-infected group, respectively.

Purification of CD4⁺ T cells and RNA isolation

Peripheral blood mononuclear cells (PBMCs) were separated and purified immediately after obtaining blood samples (30 ml whole blood from each individual) by Ficoll-Hypaque separation. Fresh CD4⁺ T cells were then obtained by positive isolation with use of microbead immunoselection (Miltenyi Biotec, Oslo, Norway). To maximize the RNA yields and minimize possible variations in gene expression profiles, the experiments were strictly followed as described in a previous study [44]. The purity of isolated CD4⁺ T cells was measured by Flow-cytometric analysis of cell markers (CD4). And it demonstrated that $98.1\% \pm 0.013$ (mean \pm SD), $98.5\% \pm 0.072$, $97.6\% \pm 0.021$ and $98.5\% \pm 0.012$ of purified CD4⁺ cells were single positive for the CD4 marker in HIV infected, HCV infected, HIV/HCV co-infected and un-infected group, respectively. For microarray analysis, RNA pooling method was used to reduce the noise and enhance reliability when many subjects were pooled without loss of individual specific information [45]. In each infection group, samples were divided into 3 subgroups for independent replicates. In each subgroup, equal amount of CD4⁺ T cells from 8 individuals were pooled together for RNA isolation and the followed gene expression profile studies. The left CD4⁺ T cells were stored at -80°C freezer for validation. RNA was isolated using RNeasy Total RNA Isolation Kit (Qiagen, Germany) and applied for microarray assays.

Microarray assays

Microarray assays were performed by using Affymetrix GeneChip Human Gene 1.0 ST Array (Affymetrix, Santa Clara, CA). A cRNA kit (The Affymetrix GeneChip Whole

Transcript Sense Target Labelling Assay) was used for complementary RNA (cRNA) generation, hybridization, and array processing. The images were acquired by the Affymetrix Scanner 3000 7G Plus and the CEL files were imported into the program Partek Genomic Suite (version 6.4, Partek Inc, St Louis, MO) and the robust multi-chip average (RMA) was normalized. Two-way ANOVA test was applied to identify differentially expressed (DE) genes (fold change >2 and adjust $p < 0.05$).

Gene set enrichment analysis

To further understand the biological meanings, Gene set enrichment analysis (GSEA) analysis (version 2.0, Broad Institute <http://www.broad.mit.edu/gsea>) was used. First, a ranked list was obtained by ranking all genes according to the correlation between their expression levels and the group distinctions using the metric signal to noise ratio. Then the association between a given gene set and the group was measured by the non-parametric running sum statistic termed the enrichment score (ES). To estimate the statistical significance of the ES, a nominal p value was calculated by permuting the genes 1,000 times. To adjust for multiple hypotheses testing, the maximum ES was normalized to account for the gene set size (NES) and the false discovery rate (FDR). The gene sets used were from Molecular Signatures Database (MsigDB), catalog C2 (version 3.0) functional sets, subcatalog canonical pathways, which include 880 gene sets from pathway databases. These gene sets were collected from online databases such as Bio-Carta, React, and KEGG (Kyoto Encyclopedia of Genes and Genomes).

Quantitative real-time RT-PCR (qRT-PCR)

RNA isolated from the CD4⁺ T cells of each participant was reverse-transcribed into cDNA using Reverse Transcription System (Promega, USA). qRT-PCR was carried out by using ABI 7500 Real-Time PCR system with Power SYBR Green Master Mix (Applied Biosystems, USA). Primer sequences are listed in Additional file 5: Table S4. The relative mRNA level of each gene was calculated as following formula: $R = 2^{CT_{(GAPDH-X)}} \times 10^3$, x represent CT value of each gene. Each RT-PCR experiment was performed three times.

Statistical analysis

Data were analyzed with SPSS version 13.0 (SPSS Inc., Chicago, IL, USA). Nonparametric test was used for pairwise comparisons. P value < 0.05 was considered as statistically significant.

Additional files

Additional file 1: Table S2. Differently expressed transcript identifiers in each comparison.

Additional file 2: Table S3. Pathways identified by gene set enrichment analysis (FDR < 0.05).

Additional file 3: Figure S1. Quantitative real time RT-PCR validations of differentially expressed genes.

Additional file 4: Table S1. Participant cohort.

Additional file 5: Table S4. Primers for Quantitative Real-time PCR Analysis.

Competing interests

The authors have declared that no competing interests exist.

Authors' contributions

MLH, ZRY, HK and JZ designed the research. JZ, LY, JL, YC, LC, JC, YS and ZL carried out all the experiments. LY and JL analyzed the data. LY wrote the manuscript. And all authors reviewed the manuscript.

Acknowledgement

This study was partially supported by National Basic Science Research Development Program of China (2013CB911302); Research Fund for Control of Infectious Diseases (RFCID, Ref. No. 09080482), the Food and Health Bureau; General Research Fund (RGC-GRF, Ref. No. 464512), The Government of the Hong Kong Special Administrative Region; Medical Scientific Project of Guangdong (A2011556) and Shenzhen Municipal Technological Project (201102106), China.

Author details

¹Stanley Ho Center for Emerging Infectious Diseases, and Li Ka Shing Institute of Health Sciences, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China. ²Shenzhen Center for Disease Control and Prevention, Shenzhen, China. ³Guangdong Provincial Institution of Public Health, Guangdong Provincial Center for Diseases Control and Prevention, Guangzhou, China. ⁴College of Life Science, Shanxi Normal University, Xi'an, China. ⁵Department of Gastroenterology, Huaxi Medical College, Sichuan University, Sichuan, China. ⁶Prince Wales Hospital, Faculty of Medicine, The Chinese University of Hong Kong, Li Ka Shing Medical Science Bldg, Hong Kong, China.

Received: 16 September 2013 Accepted: 29 January 2014

Published: 13 February 2014

References

1. Tan Y, Wei QH, Chen LJ, Chan PC, Lai WS, He ML, Kung HF, Lee SS: Molecular epidemiology of HCV mono-infection and HIV/HCV coinfection in injection drug users in Liuzhou, Southern China. *PLoS One* 2008, **3**:e3608.
2. Alter MJ: Epidemiology of viral hepatitis and HIV co-infection. *J Hepatol* 2006, **44**:S6–S9.
3. Thomas DL, Shih JW, Alter HJ, Vlahov D, Cohn S, Hoover DR, Cheung L, Nelson KE: Effect of human immunodeficiency virus on hepatitis C virus infection among injecting drug users. *J Infect Dis* 1996, **174**:690–695.
4. Liu Z, Xing WG, Zhang YH, Zhang Q, Long XS, Zhang GY, Wu H, Jiang Y: [Study on the epidemiology and HCV genotype distribution of HIV/HCV co-infection among HIV infected blood donors in China]. *Zhonghua Gan Zang Bing Za Zhi* 2006, **14**:464–465.
5. Daar ES, Lynn H, Donfield S, Gomperts E, Hilgartner MW, Hoots WK, Chernoff D, Arkin S, Wong WY, Winkler CA: Relation between HIV-1 and hepatitis C viral load in patients with hemophilia. *J Acquir Immune Defic Syndr* 2001, **26**:466–472.
6. Cribier B, Rey D, Schmitt C, Lang JM, Kim A, Stoll-Keller F: High hepatitis C viraemia and impaired antibody response in patients coinfecting with HIV. *AIDS* 1995, **9**:1131–1136.
7. Martinez-Sierra C, Arizcorreta A, Diaz F, Roldan R, Martin-Herrera L, Perez-Guzman E, Giron-Gonzalez JA: Progression of chronic hepatitis C to liver fibrosis and cirrhosis in patients coinfecting with hepatitis C virus and human immunodeficiency virus. *Clin Infect Dis* 2003, **36**:491–498.
8. Merchante N, Giron-Gonzalez JA, Gonzalez-Serrano M, Torre-Cisneros J, Garcia-Garcia JA, Arizcorreta A, Ruiz-Morales J, Cano-Lliteras P, Lozano F, Martinez-Sierra C, Macias J, Pineda JA, Grupo Andaluz para el Estudio de las Enfermedades Infecciosas: Survival and prognostic factors of HIV-infected patients with HCV-related end-stage liver disease. *AIDS* 2006, **20**:49–57.
9. d'Arminio Monforte A, Cozzi-Lepri A, Castagna A, Antinori A, De Luca A, Mussini C, Caputo SL, Arlotti M, Magnani G, Pellizzer G, Maggiolo F, Puoti M, for the Ico Foundation Study Group: Risk of developing specific AIDS-defining illnesses in patients coinfecting with HIV and hepatitis C virus with or without liver cirrhosis. *Clin Infect Dis* 2009, **49**:612–622.
10. Rockstroh JK, Mocroft A, Soriano V, Tural C, Losso MH, Horban A, Kirk O, Phillips A, Ledergerber B, Lundgren J: Influence of hepatitis C virus infection on HIV-1 disease progression and response to highly active antiretroviral therapy. *J Infect Dis* 2005, **192**:992–1002.
11. Hyrcza MD, Kovacs C, Loutfy M, Halpenny R, Heisler L, Yang S, Wilkins O, Ostrowski M, Der SD: Distinct transcriptional profiles in ex vivo CD4+ and CD8+ T cells are established early in human immunodeficiency virus type 1 infection and are characterized by a chronic interferon response as well as extensive transcriptional changes in CD8+ T cells. *J Virol* 2007, **81**:3477–3486.
12. Kovacs A, Al-Harathi L, Christensen S, Mack W, Cohen M, Landay A: CD8(+) T cell activation in women coinfecting with human immunodeficiency virus type 1 and hepatitis C virus. *J Infect Dis* 2008, **197**:1402–1407.
13. Balasubramanian A, Groopman JE, Ganju RK: Underlying pathophysiology of HCV infection in HIV-positive drug users. *J Addict Dis* 2008, **27**:75–82.
14. Ciuffreda D, Comte D, Cavassini M, Giostra E, Buhler L, Perruchoud M, Heim MH, Battegay M, Genne D, Mulhaupt B, Malinverni R, Oneta C, Bernasconi E, Monnat M, Cerny A, Chuard C, Borovicka J, Mentha G, Pascual M, Gonvers JJ, Pantaleo G, Dutoit V: Polyfunctional HCV-specific T-cell responses are associated with effective control of HCV replication. *Eur J Immunol* 2008, **38**:2665–2677.
15. Klenerman P, Hill A: T cells and viral persistence: lessons from diverse infections. *Nat Immunol* 2005, **6**:873–879.
16. Klenerman P, Kim A: HCV-HIV coinfection: simple messages from a complex disease. *PLoS Med* 2007, **4**:e240.
17. Porichis F, Kaufmann DE: HIV-specific CD4 T cells and immune control of viral replication. *Curr Opin HIV AIDS* 2011, **6**(3):174–180.
18. Wang J, Reuschel EL, Shackelford JM, Jeang L, Shivers DK, Diehl JA, Yu XF, Finkel TH: HIV-1 Vif promotes the G- to S-phase cell-cycle transition. *Blood* 2011, **117**:1260–1269.
19. Swiggard WJ, Baytop C, Yu JJ, Dai J, Li C, Schretzenmair R, Theodosopoulos T, O'Doherty U: Human immunodeficiency virus type 1 can establish latent infection in resting CD4+ T cells in the absence of activating stimuli. *J Virol* 2005, **79**:14179–14188.
20. Yoder KE, Bushman FD: Repair of gaps in retroviral DNA integration intermediates. *J Virol* 2000, **74**:11191–11200.
21. Espeseth AS, Fishel R, Hazuda D, Huang Q, Xu M, Yoder K, Zhou H: siRNA screening of a targeted library of DNA repair factors in HIV infection reveals a role for base excision repair in HIV integration. *PLoS One* 2011, **6**:e17612.
22. Bartz SR, Rogel ME, Emerman M: Human immunodeficiency virus type 1 cell cycle control: Vpr is cytostatic and mediates G2 accumulation by a mechanism which differs from DNA damage checkpoint control. *J Virol* 1996, **70**:2324–2331.
23. Izumi T, Ito K, Matsui M, Shirakawa K, Shinohara M, Nagai Y, Kawahara M, Kobayashi M, Kondoh H, Misawa N, Koyanagi Y, Uchiyama T, Takaori-Kondo A: HIV-1 viral infectivity factor interacts with TP53 to induce G2 cell cycle arrest and positively regulate viral replication. *Proc Natl Acad Sci USA* 2010, **107**:20798–20803.
24. Chang F, Re F, Sebastian S, Sazer S, Luban J: HIV-1 Vpr induces defects in mitosis, cytokinesis, nuclear structure, and centrosomes. *Mol Biol Cell* 2004, **15**:1793–1801.
25. Goh WC, Rogel ME, Kinsey CM, Michael SF, Fultz PN, Nowak MA, Hahn BH, Emerman M: HIV-1 Vpr increases viral expression by manipulation of the cell cycle: a mechanism for selection of Vpr in vivo. *Nat Med* 1998, **4**:65–71.
26. Sandberg JK, Falconer K, Gonzalez VD: Chronic immune activation in the T cell compartment of HCV/HIV-1 co-infected patients. *Virulence* 2010, **1**:177–179.
27. Chan JK, Greene WC: NF-kappaB/Rel: agonist and antagonist roles in HIV-1 latency. *Curr Opin HIV AIDS* 2011, **6**:12–18.
28. Lin W, Wu G, Li S, Weinberg EM, Kumthip K, Peng LF, Mendez-Navarro J, Chen WC, Jilg N, Zhao H, Goto K, Zhang L, Brockman MA, Schuppan D, Chung RT: HIV and HCV cooperatively promote hepatic fibrogenesis via

- induction of reactive oxygen species and NF-kappaB. *J Biol Chem* 2011, **286**:2665–2674.
29. Hiscott J, Kwon H, Genin P: **Hostile takeovers: viral appropriation of the NF-kappaB pathway.** *J Clin Invest* 2001, **107**:143–151.
 30. Liu R, Tan J, Lin Y, Jia R, Yang W, Liang C, Geng Y, Qiao W: **HIV-1 Vpr activates both canonical and noncanonical NF-kappaB pathway by enhancing the phosphorylation of IKKalpha/beta.** *Virology* 2013, **439**:47–56.
 31. de la Fuente C, Santiago F, Deng L, Eadie C, Zilberman I, Kehn K, Maddukuri A, Baylor S, Wu K, Lee CG, Pumfery A, Kashanchi F: **Gene expression profile of HIV-1 Tat expressing cells: a close interplay between proliferative and differentiation signals.** *BMC Biochem* 2002, **3**:14.
 32. Takahara Y, Takahashi M, Zhang QW, Wagatsuma H, Mori M, Tamori A, Shiomi S, Nishiguchi S: **Serial changes in expression of functionally clustered genes in progression of liver fibrosis in hepatitis C patients.** *World J Gastroenterol* 2008, **14**:2010–2022.
 33. Woodhouse SD, Narayan R, Latham S, Lee S, Antrobus R, Gangadharan B, Luo S, Schroth GP, Klenerman P, Zitzmann N: **Transcriptome sequencing, microarray, and proteomic analyses reveal cellular and metabolic impact of hepatitis C virus infection in vitro.** *Hepatology* 2010, **52**:443–453.
 34. Wu JQ, Dwyer DE, Dyer WB, Yang YH, Wang B, Saksena NK: **Genome-wide analysis of primary CD4+ and CD8+ T cell transcriptomes shows evidence for a network of enriched pathways associated with HIV disease.** *Retrovirology* 2011, **8**:18.
 35. Juno JA, Fowke KR: **Clarifying the role of G protein signaling in HIV infection: new approaches to an old question.** *AIDS Rev* 2010, **12**:164–176.
 36. Kristiansen K: **Molecular mechanisms of ligand binding, signaling, and regulation within the superfamily of G-protein-coupled receptors: molecular modeling and mutagenesis approaches to receptor structure and function.** *Pharmacol Ther* 2004, **103**:21–80.
 37. Riezu-Boj JI, Larrea E, Aldabe R, Guembe L, Casares N, Galeano E, Echeverria I, Sarobe P, Herrero I, Sangro B, Prieto J, Lasarte JJ: **Hepatitis C virus induces the expression of CCL17 and CCL22 chemokines that attract regulatory T cells to the site of infection.** *J Hepatol* 2011, **54**:422–431.
 38. Watts VJ, Neve KA: **Sensitization of adenylate cyclase by Galpha i/o-coupled receptors.** *Pharmacol Ther* 2005, **106**:405–421.
 39. Moreno-Fernandez ME, Rueda CM, Velilla PA, Rugeles MT, Chougnnet C: **cAMP during HIV infection: friend or foe?** *AIDS Res Hum Retroviruses* 2012, **28**(1):49–53.
 40. Barat C, Gilbert C, Imbeault M, Tremblay MJ: **Extracellular ATP reduces HIV-1 transfer from immature dendritic cells to CD4+ T lymphocytes.** *Retrovirology* 2008, **5**:30.
 41. Banas B, Eberle J, Schlondorff D, Luckow B: **Modulation of HIV-1 enhancer activity and virus production by cAMP.** *FEBS Lett* 2001, **509**:207–212.
 42. Bopp T, Becker C, Klein M, Klein-Hessling S, Palmethofer A, Serfling E, Heib V, Becker M, Kubach J, Schmitt S, Stoll S, Schild H, Staeger MS, Stassen M, Jonuleit H, Schmitt E: **Cyclic adenosine monophosphate is a key component of regulatory T cell-mediated suppression.** *J Exp Med* 2007, **204**:1303–1310.
 43. Hofmann B, Nishanian P, Nguyen T, Liu M, Fahey JL: **Restoration of T-cell function in HIV infection by reduction of intracellular cAMP levels with adenosine analogues.** *AIDS* 1993, **7**:659–664.
 44. Lyons PA, Koukoulaki M, Hatton A, Doggett K, Woffendin HB, Chaudhry AN, Smith KG: **Microarray analysis of human leucocyte subsets: the advantages of positive selection and rapid purification.** *BMC Genomics* 2007, **8**:64.
 45. Peng X, Wood CL, Blalock EM, Chen KC, Landfield PW, Stromberg AJ: **Statistical implications of pooling RNA samples for microarray experiments.** *BMC Bioinforma* 2003, **4**:26.

doi:10.1186/1743-422X-11-27

Cite this article as: Yi et al.: Gene expression profiling of CD4⁺ T cells in treatment-naïve HIV, HCV mono- or co-infected Chinese. *Virology Journal* 2014 **11**:27.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

