

SHORT REPORT

Open Access

Hepatitis G Virus associated aplastic anemia: A recent case from Pakistan

Shahida Amjad Riaz Shah¹, Muhammad Idrees^{2*}, Abrar Hussain²

Abstract

Background: Aplastic anemia (AA) is a serious and rare disorder characterized by a hypocellular bone marrow. Hepatitis associated aplastic anemia (HAAA) is a variant of aplastic anemia in which aplastic anemia follows an acute attack of hepatitis. Several reports have noted an association between HGV and hepatitis-associated aplastic anemia besides other hepatitis causing viruses.

Case presentation: A female girl of age 11 year with a history of loose motion for one month, vomiting for last 15 days and poor oral intake for last few days is reported here. The physical examination presents fever, pallor whereas bleeding, hepatomegaly, Splenomegaly and bruising were absent, abdominal ultrasonography confirmed the absence of hepatomegaly, Splenomegaly and lymphadenopathy. The laboratory investigation parameters were: haemoglobin 6.2 g/L, total leucocytes count 1.51, neutrophils 0.47%, absolute reticulocyte count 0.5%, Monocytes 0.16%, red cell count 3.2 mil/uL, Picked cell volume (PCV) 30.13%, Mean Corpuscular Volume (MCV) 78 fL, Mean Corpuscular Hemoglobin (MCH) 26.3 pg. The liver enzymes were alanine aminotransferase (ALT) 98 IU/L, aspartate aminotransferase (AST) 114 IU/L. Serologic and molecular tests for hepatitis A, B, C, D, E, TTV, B19 were negative, whereas HGV RNA PCR test was found positive for hepatitis G virus. The bone marrow aspirate and trephine biopsy examination revealed hypo- cellularity, erythropoiesis, myelopoiesis and megakaryopoiesis.

Conclusion: HAAA is an uncommon but severe condition, which may occur following idiopathic cases of acute hepatitis. Our finding suggests the involvement of HGV in the development of aplastic anemia. In patients presenting with pancytopenia after an episode of acute hepatitis, the definitive diagnosis should be considered and confirmed by RT-PCR and if possible by bone marrow biopsy.

Introduction

Hepatitis G virus was reported first time has a non-A-E hepatitis and placed as flavivirus [1]. The induction of this new agent in the family of Hepatitis has attracted significant attention because of its etiology [2]. Hepatitis G virus has been marked as a cause of non-A through E acute viral hepatitis and sharp liver failure. Aplastic anemia complicating hepatitis is an uncommon but well recognized phenomenon. Hepatitis associated aplastic anemia is a severe disorder with a high mortality (85%) [3]. Hepatitis associated aplastic anemia (HAAA) is a deviation of aplastic anemia in which aplastic anemia follows an acute attack of hepatitis. The marrow failure can be severe and is

usually lethal if untreated. Lorenz and Quazier has documented first time HAAA in two case back in 1955 [4], by 1975 more than 193 cases had been reported [5]. Adil et al has reported, severe aplastic anaemia (SAA) 51.4%, very severe (VSAA) in 16.7% of 144 patients of aplastic anemia cases [6]. A number of reports have mentioned alliance between HGV and HAAA [7-10]. Number of HAAA cases with a history of multiple blood transfusions has been reported [11,12]. Crespo et al has documented a case of 24 year old man have community acquired HGV that later progress into severe aplastic anemia, point out HGV for both hepatitis and aplastic anaemia. However greater number of serum samples are needed to prove the association of hepatitis G virus and aplastic anaemia [13]. Moatter et al. has reported 5/43 patients of haemodialysis with raised liver enzyme, reduced platelet count and 21/100 patients of polytransfused b- thalassemia major children infected with HGV RNA form Pakistan [14,15].

* Correspondence: idreeskhan96@yahoo.com

²Division of Molecular Virology, National Centre of Excellence in Molecular Biology, 87-West Canal Bank Road Thokar Niaz Baig Lahore-53700, University of the Punjab Lahore, Pakistan

Full list of author information is available at the end of the article

Case Presentation

In this study well characterized samples of 93 aplastic anaemia patients before blood transfusion were included. These characteristics include history, physical examination, haematological investigation, bone marrow aspirate and trephine biopsy examination, liver function test (LFTs), renal parameters, viral profile and abdominal ultrasonography. The diagnosis of HA-aplastic anemia was made on the basis of hepatitis (elevated serum aminotransferase enzymes, jaundice, absolute neutrophils counts, platelet counts and reticulocytes. All the 93 samples were checked for serological marker of HAV, HBV, HCV, HDV, HEV, HGV, TTV and B19. One of 93 samples from patients with HA-aplastic anemia has hepatitis G associated aplastic anaemia with positive HGV RNA.

A female girl of age 11 year is reported here. The patient had a history of loose motion for one month, vomiting for last 15 days and poor oral intake for last few days. The physical examination presents fever, pallor whereas bleeding, hepatomegaly, Splenomegaly and bruising were absent, abdominal ultrasonography confirmed the absence of hepatomegaly, Splenomegaly and lymphadenopathy. The laboratory investigation parameters were: haemoglobin 6.2 g/L, total leucocytes count 1.51, neutrophils 0.47%, absolute reticulocyte count 0.5%, Monocytes 0.16%, red cell count 3.2 mil/uL, Picked cell volume (PCV) 30.13%, Mean Corpuscular Volume (MCV) 78 fL, Mean Corpuscular Hemoglobin (MCH) 26.3 pg. The liver enzymes were alanine aminotransferase (ALT) 98 IU/L, aspartate aminotransferase (AST) 114 IU/L. Serologic and molecular tests for hepatitis A, B, C, D, E, TTV, B19 were negative, whereas HGV RNA PCR test was found positive for hepatitis G virus. The bone marrow aspirate and trephine biopsy examination revealed hypo-cellularity, erythropoiesis, myelopoiesis and megakaryopoiesis.

Discussion

Flaviviruses belong to enveloped viruses with a single positive sense RNA about 10 kb. Hepatitis G virus medium of transmission mostly through blood. The possible role of hepatitis G virus infection in the pathogenesis of rare non-liver disease has been suggested but need to be recognized. By some unknown reason aplastic anemia some time preceded by hepatitis. Few reports have analysed the role of HGV in the development of HAAA [8,9,11]. Crespo et al in print a case through negative serological markers for HAV, HBV, HCV, HEV, hypoplastic marrow low platelet and white cell counts but detected HGV-RNA before any blood transfusion [13]. Byrnes et al has described hepatitis G Virus positive case of 26 year old man before the use of medication, blood transfusion or intravenous drug abuse [9]. Zaidi et al,

reported a 19 year male before blood transfusion with positive HGV by RT-PCR and suggested that in the absence of any other clinical manifestations the possible infectious agent may be HGV for hepatitis G virus associated aplastic anaemia [8]. In the list of studies of Hepatitis G Virus associated aplastic anaemia before blood transfusion we report a case of 11 years female girls.

Whereas some reports confirm the presence of HGV viraemia in the patient aplastic anaemia after blood transfusion [8,9,11,12,16-18]. But however concluded that HGV viraemia is frequent in patients with aplastic anaemia [19]. Kiem et al reported similar finding after logistic regression analysis, that HGV RNA in transfused patients was 5.9 times higher compared to untransfused patients ($P = 0.001$). This implicates transfusion as major source of HGV with aplastic anaemia [18].

The published literature point out that studies must be performed on many more aplastic anaemia patients prior to blood transfusion [20]. However, to find such patients in large number are not normally available to study, so far individual cases are reported. The ideal case regarding Hepatitis G associated aplastic anaemia are pre blood transfusion.

Conclusion

In conclusion, HAAA is an uncommon but severe condition, which may occur following idiopathic cases of acute hepatitis. Our finding suggests the involvement of HGV in the development of aplastic anemia. In patients presenting with pancytopenia after an episode of acute hepatitis, the definitive diagnosis should be considered and confirmed by RT-PCR and if possible by bone marrow biopsy.

Consent

Written informed consent was obtained from the patients a copy of which is available for review by Editor-in-Chief of this journal.

Author details

¹Hematology Department, Sheikh Zayed Medical Complex, Lahore Pakistan.
²Division of Molecular Virology, National Centre of Excellence in Molecular Biology, 87-West Canal Bank Road Thokar Niaz Baig Lahore-53700, University of the Punjab Lahore, Pakistan.

Authors' contributions

SARS aided in acquisition of data that was included in this case report and drafted the manuscript. MI aided in acquisition and interpretation of the data analysed. AH helped in statistical analysis of the data and in editing of this manuscript. All authors have read and approved the final version of this manuscript.

Authors Information

SARS: Consultant Hematologist, Sheikh Zayed Medical Complex. MI: Consultant Molecular Virologist, Ministry of Education. AH: Assistant Professor and PhD Research Scholar.

Competing interests

The authors declare that they have no competing interests.

Received: 29 December 2010 Accepted: 21 January 2011

Published: 21 January 2011

References

1. Simons JN, Pilot-Matias TJ, Leary TP, Dawson GJ, Desai SM, Schlauder GG, et al: **Identification of two flavivirus-like genomes in the GB hepatitis agent.** *Proc Natl Acad Sci USA* 1995, **92**(8):3401-5.
2. Reshetnyak VI, Karlovich TI, Ilchenko LU: **Hepatitis G virus.** *World J Gastroenterol* 2008, **14**(30):4725-34.
3. Fatemeh Farahmand GHF, Touran Shahraki, Gholam Reza Khatami, Mina Izadiar GAR: **Hepatitis-Associated Aplastic Anemia.** *Govaresh* 2005, **10**(1):54-8.
4. Lorenz EQK: **Panmyelopathy following epidemic hepatitis.** *Wien Med Wochenschr* 1955, **8**(105(1)):19-22.
5. Hagler L, Pastore RA, Bergin JJ, Wrensch MR: **Aplastic anemia following viral hepatitis: report of two fatal cases and literature review.** *Medicine, Baltimore* 1975, **54**(2):139-64.
6. Adil SN, Burney IA, Kakepoto GN, Khurshid M: **Epidemiological features of aplastic anaemia in Pakistan.** *J Pak Med Assoc* 2001, **51**(12):443-5.
7. Hibbs JR, Frickhofen N, Rosenfeld SJ, Feinstone SM, Kojima S, Bacigalupo A, et al: **Aplastic anemia and viral hepatitis. Non-A, Non-B, Non-C?** *JAMA* 1992, **267**(15):2051-4.
8. Zaidi Y, Chapman CS, Myint S: **Aplastic anaemia after HGV infection.** *Lancet* 1996, **348**(9025):471-2.
9. Byrnes JJ, Banks AT, Piatak M Jr, Kim JP: **Hepatitis G-associated aplastic anaemia.** *Lancet* 1996, **348**(9025):472.
10. Linnen J, Wages J Jr, Zhang-Keck ZY, Fry KE, Krawczynski KZ, Alter H, et al: **Molecular cloning and disease association of hepatitis G virus: a transfusion-transmissible agent.** *Science* 1996, **271**(5248):505-8.
11. Brown KE, Tisdale J, Barrett AJ, Dunbar CE, Young NS: **Hepatitis-associated aplastic anemia.** *N Engl J Med* 1997, **336**(15):1059-64.
12. Kiem HP, Storb R, McDonald GB: **Hepatitis-associated aplastic anemia.** *N Engl J Med* 1997, **337**(6):424-5.
13. Crespo JdlHB, Rivero M, Lozano JL, Fábrega E, Pons-Romero F: **Hepatitis G virus infection as a possible causative agent of community-acquired hepatitis and associated aplastic anaemia.** *Postgrad Med J* 1999, **88**(75):159-60.
14. Moatter T, Tariq SA, Fisher-Hoch S, Haroon S, McCormick JB: **Hepatitis G virus in Karachi, Pakistan.** *Lancet* 1996, **348**(9037):1318-9.
15. Moatter T, Adil S, Haroon S, Azeemuddin S, Hassan F, Khurshid M: **Prevalence of hepatitis G virus in Pakistani children with transfusion dependent beta- thalassemia major.** *Indian J Pathol Microbiol* 1999, **42**(4):475-82.
16. Kiem HP, Myerson D, Storb R, McDonald GB, Spurgeon CL, Leisenring W: **Prevalence of hepatitis G virus in patients with aplastic anemia.** *Blood* 1997, **90**(3):1335-6.
17. Alter HJ, Nakatsuji Y, Melpolder J, Wages J, Wesley R, Shih JW, et al: **The incidence of transfusion-associated hepatitis G virus infection and its relation to liver disease.** *N Engl J Med* 1997, **336**(11):747-54.
18. Moriyama K, Okamura T, Nakano S: **Hepatitis GB virus C genome in the serum of aplastic anaemia patients receiving frequent blood transfusions.** *Br J Haematol* 1997, **96**(4):864-7.
19. Brown KE, Wong S, Young NS: **Prevalence of GBV-C/HGV, a novel 'hepatitis' virus, in patients with aplastic anaemia.** *Br J Haematol* 1997, **97**(2):492-6.
20. Robaczewska M, Cova L, Podhajska AJ, Falkiewicz B: **Hepatitis G virus: molecular organization, methods of detection, prevalence, and disease association.** *Int J Infect Dis* 1999, **3**(4):220-33.

doi:10.1186/1743-422X-8-30

Cite this article as: Riaz Shah et al.: **Hepatitis G Virus associated aplastic anemia: A recent case from Pakistan.** *Virology Journal* 2011 **8**:30.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

