

RESEARCH

Open Access

Identification of B cell epitopes reactive to human papillomavirus type-16L1- derived peptides

Akimasa Fukui¹, Satoko Matsueda², Kouichiro Kawano^{1,4*}, Naotake Tsuda¹, Nobukazu Komatsu², Shigeki Shichijo², Tetsuro Sasada², Satoshi Hattori³, Kimio Ushijima¹, Kyogo Itoh² and Toshiharu Kamura¹

Abstract

Background: Persistent infection of human papillomavirus (HPV) types 16 and 18 causes cervical cancer. To better understand immune responses to the prophylactic vaccine, HPV 16/18 L1 virus-like particles (HPV-VLPs), we investigated B cell epitopes of HPV16 L1-derived peptides.

Methods: Sera from mice immunized with HPV-16/18 L1 VLPs were analyzed for their IgG titers against 10 different HPV16 L1-derived peptides (20-mer) that contain human leukocyte antigen (HLA)-class I A-2, A-24 and class II DR.

Results: One 20-mer peptide at positions 300 to 319 was identified as a common B cell epitope in both Balb/c (H-2^d) and C57BL/6 (H-2^b) mice. Mapping analysis showed that the 10-amino-acid sequence at positions 304 to 313 was an immunogenic portion. It is of note that the binding capability of this 10-mer peptide to the HLA-A2 and HLA-A24 molecules was confirmed by the HLA class I stabilization assay. In addition, one unique 20-mer was determined as a B cell epitope in each strain.

Conclusions: These results might provide new information for better understanding of immune responses to HPV 16 L1.

Keywords: Human papillomavirus, Prophylactic vaccine, Anti-peptide antibody, Virus-like particles

Background

Cervical cancer is the second most prevalent cancer in women worldwide. HPV 16 is the most common type associated with cervical cancer [1]. HPV-16/18L1 virus-like particles (HPV-VLPs), which induce neutralizing antibody responses, have been used as a prophylactic vaccine with great success [2,3]. Although the preventive effect of the HPV-VLPs vaccine has been reported to last up to 7.3 years [4], the durability is unclear at the present time, either for the entire vaccinated population or for individuals. This hurdle could be in part overcome if predictable biomarkers were identified. One of the biomarkers could be based on the measurement of specific humoral immune responses to the vaccine. However, little information is presently available with regard to humoral responses against the HPV-VLPs, primarily

because of the limited availability of the assay reagents [5,6]. The main aim of this study was to better understand humoral immune responses to HPV16 L1-derived peptides in an animal model.

Results

Detection of IgG antibodies in serum of Balb/c mice


Serum of the Balb/c mice was obtained before immunization, and 3, 5, 8, 11, 14 weeks after the first immunization. Each group consisted of 6 mice, and serum from each mouse was independently measured for IgG level. A 100-fold dilution of samples was used to determine the levels of IgG reactive to each of 10 different HPV16 L1-derived peptides (20-mer), and the results were given in fluorescent intensity units (FIU) (Figure 1). Representative results of a kinetic study showed that IgG against peptide 4 and peptide 6, but not any others, became detectable at 3 weeks and reached a maximum at 5 weeks ($P < 0.05$) followed by decline thereafter until 11 weeks. IgG levels were somewhat increased again at 14 weeks, which might be in part a reflection of the

* Correspondence: kawano_kouichirou@kurume-u.ac.jp

¹Department of Obstetrics and Gynecology, Kurume University School of Medicine, Kurume, Japan

⁴Department of Obstetrics and Gynecology, Kurume University School of Medicine, 67 Asahi-machi, Kurume, Fukuoka 830-0011, Japan

Full list of author information is available at the end of the article



second immunization at 3 weeks after the first injection. A dim level of IgG against peptide 8 was detectable at 8 and 11 weeks, but the levels were not significant.

Detection of IgG antibodies in serum of C57BL/6 N (H-2^b) mice

We then measured IgG levels in serum from the vaccinated C57BL/6N (H-2^b) mice before and 5 weeks after the immunization to address any difference between Balb/c and C57BL/6N mice. As a result, IgG against both peptide 6 and peptide 8, but not any others, were detected at 5 weeks ($P < 0.05$) (Figure 2).

Epitope mapping of peptide

These results indicated that peptide 6 is the major common B cell epitope shared by the two strains. Subsequently, epitope mapping of peptide 6 was conducted with eight different 10-mers (Figure 3). Five amino acid sequences from the HPV16 L1-derived sequence at position 295 to 299 were added to the N-terminal of the first 10-mer, and those at position 320 to 324 were also added to the C-terminal of the eighth 10-mer. Each of these eight peptides shared seven amino acid sequences with each other. As a result, one 10-mer at amino acid positions of 304–313 of HPV16 L1, aqifnkpywl, was determined to be an immunogenic portion using the antibodies (Figure 3). Then, we addressed the reactivity of the 9-mer peptide (qifnkpywl) at position 305 to 313, which had binding motifs to HLA-A2 and HLA-A24, to immunized sera. A modest level of reactivity to this 9-mer was observed (Figure 3).


Binding capability of the immunogenic B cell epitope to HLA class I molecules

The identified immunogenic B cell epitope contains binding motifs to the HLA-A2 and HLA-A24 molecules. Therefore, we examined whether they actually bind to the HLA-A2- and HLA-A24 molecules by the HLA class I stabilization assay with TAP-deficient cell lines RMA-S/A2 and RMA-S/A24. As illustrated in Table 1, both of the 9-mer (qifnkpywl) and 10-mer (aqifnkpywl) peptides showed substantial binding capability to HLA-A2 and HLA-A24.

Discussion

We analyzed immune responses to the 10 different HPV-VLP L1-derived peptides (20-mers) that had binding motifs to both HLA-class I (A2 or A24) and HLA-class II (DR) in animal model. We used BALB/c and C57BL/6N mice that have been regarded as Th2- and Th1-skewed strains, respectively, and widely known to express different immune responses in normal and pathological states [7]. When we examined humoral immune responses to the 10 different HPV-VLP L1-derived peptides in the Th2-skewed BALB/c mice, the levels of IgG to the peptide 4 and peptide 6 were clearly elevated in the sera after immunization with HPV-VLPs. We also used the Th1-skewed C57BL/6N mice to examine whether the selection of mouse strains tested has considerable and variable impacts on humoral immune responses to HPV-VLPs. In the C57BL/6N mice, one of the identified peptides, peptide 6, but not another one (peptide 4), was also immunogenic, suggesting that the peptide 6 is a major common B cell epitope peptide in mice. Notably, our preliminary study has shown that the peptide 6 is also immunogenic in humans who are vaccinated with HPV-VLPs (data not shown), suggesting that the peptide 6 is a common immunogenic B cell epitope shared between mice and humans. A dim level of IgG against peptide 8 was detected in Balb/c mice; thus, peptide 8 might also contain a common epitopic portion. This issue remains to be studied further.

Mapping analysis of peptide 6 showed 10-mer (aqifnkpywl) at positions of 304 to 313 of HPV16 L1 as an immunogenic portion. In addition, the 9-mer peptide (qifnkpywl) at position 305 to 313 with binding motifs to HLA-A2 and -A24 molecules was also recognized by immunized sera. It is of note that both these 9-mer and 10-mer peptides showed substantial binding capability to HLA-A2 and HLA-A24 molecules, the two dominant HLA-class I A types among Japanese and other ethnics [8]. The peptide 6 at positions 300 to 319 of HPV16 L1 also contains binding motifs to HLA-class-II DR 1, 3, and 4 molecules, the dominant molecules expressed in Japanese and other ethnics [9]. Therefore, the peptide 6 and its fragment might be one of the appropriate



candidate antigenic sequences for monitoring immune responses to HPV-VLP after vaccination both in animal model and humans. This hypothesis is now under investigation in humans.

Whether the vaccine-induced IgG to the peptide 6 or 8 possesses biological activity to either neutralize HPV infection or facilitate the prophylactic effect of the HPV 16/18L1 vaccine needs to be studied in near future. In addition, since we have known that the same epitopes

are often recognized by both B cells and T cells [10], T cell responses to the peptide 6 and its fragment may also be of great interest.

Conclusions

In summary, one 20-mer peptide at positions 300 to 319 was identified as a common B cell epitope in both Balb/c (H-2^d) and C57BL/6N (H-2^b) mice. These results




Table 1 HLA-binding capability of the B cell epitope derived from HPV16 L1

Peptide	Sequence	Binding capability (%)*	
		HLA-A2	HLA-A24
9-mer	QIFNKPYWL	66.8	40.3
10-mer	AQIFNKPYWL	74.9	51.7
Flu M1	GILGFVFTL	116.7	ND
EBV	TYGPVFMCL	ND	170.6
K-Ras	KLWVG AGGV	3.3	3.8

*HLA-A2 or HLA-A24-binding capability was estimated by increase in mean fluorescence intensity (MFI), determined by flowcytometry after staining of RMA-S/A2 or RMA-S/A24 cells with anti-HLA-A2 or anti-HLA-A24 mAb, as follows; MFI increase (%) = (MFI with a given peptide - MFI without peptide) / (MFI without peptide) X 100. As positive controls, an HLA-A2-binding peptide derived from influenza virus M1 (Flu M1) or an HLA-A24-binding peptide derived from Epstein-Barr virus LMP2 (EBV) was used. As a negative control, a peptide derived from oncogene K-ras was used. ND, not determined.

might provide new information for better understanding of immune responses to HPV 16L1.

Materials and methods

Immunization of mice

Female Balb/c (H-2^d) and C57BL/6N (H-2^b) mice used in this study were 6 weeks of age and were maintained in a pathogen-free environment. Balb/c mice (H-2^d) were mainly provided for the study because they have been regarded as a Th2- skewed strain [7] and the binding motifs of the peptides to the H-2^d class I-A molecules were similar to those of human leukocyte antigen (HLA) class I-A2402, which are expressed in 60% of Japanese [11]. We administered the bivalent HPV-16/18 virus-like particle AS04 vaccine, containing 2 µg each of HPV-16 and HPV-18L1 VLPs with AS04 adjuvant that contained 50 µg aluminum hydroxide and 5 µg 3-deacylated monophosphoryl lipid A. Mice were immunized twice (day 1 and day 21) into the muscle at the

thigh with HPV16/18 VLP AS04 vaccine based on the report by Didierlaurent et al. [12].

All experiments with live animals were performed under the institutional guidelines of the animal experiment in Kurume University after the approval by the Committee of Animal Experiment in Kurume University (Approval Number: 2010-107-1).

Collection of serum samples

The blood of each mouse was collected from the orbital sinus. Blood collection was performed before vaccination, and 3, 5, 8, 11, and 14 weeks after the first immunization in Balb/C (n=6 in each group). In C57BL/6 mice, blood was collected before (n=9) and 5 weeks after the first immunization (n=9). Collected blood was allowed to coagulate by keeping it at room temperature for 1 h followed by centrifugation and collection. Serum from each mouse was sealed and stored at -80°C until use. Measurement of IgG titers was conducted simultaneously to avoid possible *in vitro* biases.

Peptides

Ten different HPV16 L1-derived peptides (20-mer) with binding motifs to both HLA-class I (A2 or A24) and HLA-class II (DR) were selected by the web software (MULTIPRED) (Table 2). This choice was based on consideration of future applications to the human immune system. For epitope mapping, 8 different 10-mer and one 9-mer peptides were selected from the 20-mer peptide 6. These peptides were purchased from Greiner Bio-One (Thermo Fisher Scientific, Ulm, Germany). Each peptide was dissolved in dimethyl sulfoxide (DMSO), stored at -80°C.

Preparation of xMAP beads

The xMAP carboxylate beads and Luminex system platform were obtained from Luminex Corp. (Austin, TX) as

Table 2 HPV16 L1-derived peptides used in this study and their binding motifs to HLA-A2 and -A24

	DR			A2			4A2		
	position	Amino acid sequence	Score	position	Amino acid sequence	Score	position	Amino acid sequence	Score
Peptide 1	54-73	KPNNNKILVPKVSGLQYRVF	30	60-68	ILVPKVSGL	30	59-68	KILVPKVSGL	14
Peptide 2	392-422	HSMNSTILEDWNFGLQPPPGG	23	398-406	ILEDWNFGL	23	397-406	TILEDWNFGL	16
Peptide 3	62-81	VPKVSGLQYRVFRIHLPDPN	22	67-75	GLQYRVFRI	22	66-75	SGLQYRVFRI	24
Peptide 4	112-131	PLGVGISGHPLLKDDTEN	22	118-126	SGHPLLNKL	22	117-126	ISGHPLLNKL	12
Peptide 5	243-262	GDSLFFYLRRREQMFVRHLFN	22	249-257	YLRRREQMFV	22	248-257	FYLRRREQMFV	12
Peptide 6	300-319	VTSDAQIFNKPYWLQRAQGH	21	305-313	QIFNKPYWL	21	305-313	QIFNKPYWL	12
Peptide 7	144-162	RECISMDYKQTQLCLIGCK	20	148-156	SMDYKQTQL	20	148-156	SMDYKQTQL	11
Peptide 8	293-312	PTPSGSMVTSDAQIFNKPYW	20	298-306	SMVTSDAQI	20	298-306	SMVTSDAQI	10
Peptide 9	384-403	TADVMTYIHSMNSTILEDWN	20	390-399	YIHSMNSTIL	20	389-398	TYIHSMNSTI	23
Peptide 10	152-171	KQTQLCLIGCKPPIGEHWG	23	157-165	CLIGCKPPI	23	156-165	LCLIGCKPPI	12

Anchor residues for HLA class I are shown in boldface.

reported previously [13]. The 96-well filter plates (MABVN12) and vacuum manifold apparatus (MAVM 09601) were from Millipore Corp. (Bedford, MA). Biotinylated goat anti mouse IgG (gamma chain-specific) (SouthernBiotech, AL) was purchased from Vector Laboratories Inc. (Burlingame, CA). Streptavidin-PE (S-866) was purchased from Molecular Probes (Eugene, OR). 1-Ethyl-3-[3-dimethylaminopropyl] carbodiimide hydrochloride (EDC, 22980) was obtained from PIERCE (Rockford, IL). Peptides were coupled to xMAP beads according to the modified manufacturer's instructions as reported previously [13]. In brief, 100 μ of xMAP beads were washed with 0.1 M MES buffer, pH 7.0, followed by mixing with 100 μ l of peptide (1 mg/ml in 0.1 M MES buffer, pH 7.0). The peptide-loaded beads were incubated with EDC (1 mg/ml) at room temperature for 30 min in darkness, and then incubated twice more under the same conditions, after which the beads were washed with 0.05% Tween 20-PBS. Finally, the beads were treated with 2-aminoethanol for 15 min at room temperature in darkness, then washed twice and re-suspended with 1 ml of 0.05% NaN₃ in Block-Ace.

Anti-peptide antibody measurement by multiplexed bead-based Luminex assay

Blood samples were obtained from each of the mice at each scheduled point. Peptide-specific IgG levels in serum were measured by flowmetry assay using the Luminex system as reported previously [13]. In brief, serum was incubated with 100 μ l of the peptide-coded beads for 1.5 hours at room temperature in a 96-well filter plate on a plate shaker. After incubation, the plate was washed using a vacuum manifold apparatus and incubated with 100 μ l of biotinylated goat anti mouse IgG (gamma chain-specific) for 1 hour at room temperature on a plate shaker. The plate was then washed, 100 μ l of streptavidin-PE was added to the wells, and the plate was incubated for 40 min at room temperature on a plate shaker. The bound beads were washed three times followed by the addition of 100 μ l of Tween 20-PBS into each well, and the plate was placed for 3 min on a plate shaker.

HLA class I stabilization assay

The actual binding of the peptides to HLA-A2 or HLA-A24 molecules was evaluated by MHC class I stabilization assay with the TAP2-deficient RMA-S cells stably transfected with the HLA-A*0201 gene (RMA-S/A2) or with the HLA-A*2402/K^b gene (RMA-S/A24), according to a previously reported method with several modifications [14]. Briefly, RMA-S/A2 or RMA-S/A24 cells (5x10⁵ cells per well in a 24-well plate) were cultured for 18 hours at 26°C in 1 ml of RPMI 1640 medium (Invitrogen Inc, Carlsbad, CA) containing 10% FBS (MP Biologicals, Solon, OH) in the presence of

synthetic peptides (25 μ g/ml) and β 2-microglobulin (2 μ g/ml; Fitzgerald Industries International, Acton, MA). After washing, the cells were cultured for 3 hours at 37°C, and then stained with anti-HLA-A2 mAb (BB7.2; BD Bioscience, San Jose, CA) or anti-HLA-A24 mAb (One Lambda, Inc. Canoga Park, CA), followed by incubation with PE-conjugated rabbit anti-mouse IgG Ab (MP Biomedicals, Solon, OH). After washing, the cells were suspended with 1 ml of PBS containing 1% formaldehyde, and analyzed with FACSCanto (BD Bioscience). The binding capability of each peptide to HLA-A2 or HLA-A24 molecules was evaluated by the increase in mean fluorescence intensity (MFI) assessed by flow cytometry, as follows: MFI increase (%) = (MFI with a given peptide - MFI without peptide)/(MFI without peptide) X 100. As positive controls, an HLA-A2-binding peptide derived from influenza virus M1 (Flu M1, GILGFVFTL) or an HLA-A24-binding peptide derived from Epstein-Barr virus LMP2 (EBV, TYGPVFMCL) was used. As a negative control, a peptide derived from oncogene K-ras (KLVVVG AGGV) was used.

Statistics

The statistical significance of the data was determined using Friedman's test and the Wilcoxon rank sum test. P-values less than 0.05 were considered statistically significant.

Abbreviations

HPV: Human papillomavirus; VLPs: Virus-like particles; HLA: Human leukocyte antigen; FIU: Fluorescent intensity units.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

A.F. performed most of the experiments and involved in manuscript preparation. T.S. and N.K. coordinated laboratory manipulation. KK proposed the hypothesis of this study edited the manuscript. K.K., N.T., K.U., K.I. and T.K. designed this study. S.S. and A.F. designed the peptides analyzed. A.F., S.M. gave vaccination to mice and obtained blood sample, A.F., S.M., N.K. and T.S. measured the antibody. A.F., T.S., S. H., and K.I. analyzed the data. KI and TK are the project leaders and were involved in project design, manipulation, data analysis and finalization of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported in part by the Supporting Fund of Obstetrics and Gynecology, Kurume University, by the grant from Regional Innovation Cluster Program of the Ministry of Education, Culture, Sports, Science and Technology of Japan, and by the the grant from Sendai-Kousei Hospital.

Author details

¹Department of Obstetrics and Gynecology, Kurume University School of Medicine, Kurume, Japan. ²Department of Immunology and Immunotherapy, Kurume University School of Medicine, Kurume, Japan. ³Biostatistics Center, Kurume University, Kurume, Japan. ⁴Department of Obstetrics and Gynecology, Kurume University School of Medicine, 67 Asahi-machi, Kurume, Fukuoka 830-0011, Japan.

Received: 22 February 2012 Accepted: 11 September 2012
Published: 14 September 2012

References

1. Muñoz N, Bosch FX, de Sanjosé S, Herrero R, Castellsagué X, Shah KV, Snijders PJ, Meijer CJ: **International Agency for Research on Cancer Multicenter Cervical Cancer Study Group: Epidemiologic classification of human papillomavirus types associated with cervical cancer.** *N Engl J Med* 2003, **348**:518–527.
2. Villa LL, Costa RL, Petta CA, Andrade RP, Ault KA, Giuliano AR, Wheeler CM, Koutsky LA, Malm C, Lehtinen M, Skjeldestad FE, Olsson SE, Steinwall M, Brown DR, Kurman RJ, Ronnett BM, Stoler MH, Ferenczy A, Harper DM, Tamms GM, Yu J, Lupinacci L, Railkar R, Taddeo FJ, Jansen KU, Esser MT, Sings HL, Saah AJ, Barr E: **Prophylactic quadrivalent human papillomavirus (types 6, 11, 16, and 18) L1 virus-like particle vaccine in young women: a randomised double-blind placebo-controlled multicentre phase II efficacy trial.** *Lancet Oncol* 2005, **6**:271–278.
3. Harper DM, Franco EL, Wheeler C, Ferris DG, Jenkins D, Schuind A, Zahaf T, Innis B, Naud P, De Carvalho NS, Roteli-Martins CM, Teixeira J, Blatter MM, Korn AP, Quint W, Dubin G, GlaxoSmithKline HPV Vaccine Study Group: **Efficacy of a bivalent L1 virus-like particle vaccine in prevention of infection with human papillomavirus types 16 and 18 in young women: a randomised controlled trial.** *Lancet* 2004, **364**:1757–1765.
4. De Carvalho N, Teixeira J, Roteli-Martins CM, Naud P, De Borba P, Zahaf T, Sanchez N, Schuind A: **Sustained efficacy and immunogenicity of the HPV-16/18 AS04-adjuvanted vaccine up to 7.3 years in young adult women.** *Vaccine* 2010, **28**:6247–6255.
5. Joura EA, Kjaer SK, Wheeler CM, Sigurdsson K, Iverson OE, Hernandez-Ailta M: **HPV antibody levels and clinical efficacy following administration of a prophylactic quadrivalent HPV vaccine.** *Vaccine* 2008, **26**:6844–6851.
6. Ryding J, Dahlberg L, Wallen-Ohman M, Dilner J: **Deletion of a major epitope of human papillomavirus type 16 virus-like particles.** *J Gen Virol* 2007, **88**:792–802.
7. Tacchini-Cottier F, Weinkopff T, Launois P: **Does T Helper Differentiation Correlate with Resistance or Susceptibility to Infection with L. major? Some Insights From the Murine Model.** *Front Immunol* 2012, **3**:32.
8. Rammensee GH, Fried T, Stevanovic S: **MHC ligands and peptide motifs: first listing.** *Immunogenetics* 1995, **41**:178–228.
9. Guang Lan Z, Khan AM, Srinivasan KN, August JT, Bruscia V: **MULTIPRED: a computational system for prediction of promiscuous HLA binding peptides.** *Nucleic Acids Res* 2005, **33**:172–179.
10. Mine T, Sato Y, Noguchi M, Sasatomi T, Gouhara R, Tsuda N, Tanaka S, Shomura H, Katagiri K, Rikimaru T, Shichijo S, Kamura T, Hashimoto T, Shirouzu K, Yamada A, Todo S, Itoh K, Yamana H: **Humoral responses to peptides correlate with overall survival in advanced cancer patients vaccinated with peptides based on pre-existing peptide-specific cellular responses.** *Clin Cancer Res* 2004, **10**(3):929–937.
11. Imanishi T, Akazawa T, Kimura A: **Allele and haplotype frequencies for HLA and complement loci in various ethnic groups.** In *HLA 1991*, Volume 1. Edited by Tsuji K, Aizawa M, Sasazuki T. Oxford: Oxford Scientific Publications; 1992:1065–1220.
12. Didierlaurent AM, Morel S, Lockman L, Giannini SL, Bisteau M, Carlsen H, Kielland A, Vosters O, Vanderheyde N, Schiavetti F, Larocque D, Van Mechelen M, Garçon N: **AS04, an aluminum salt- and TLR4 agonist-based adjuvant system, induces a transient localized innate immune response leading to enhanced adaptive immunity.** *J Immunol* 2009, **183**(10):6186–6197.
13. Komatsu N, Shichijo S, Nakagawa M, Itoh K: **New multiplexed flow cytometric assay to measure anti-peptide antibody: a novel tool for monitoring immune responses to peptides used for immunization.** *Scand J Clin Lab Invest* 2004, **64**:535–546.
14. Nakao M, Shichijo S, Imaizumi T, Inoue Y, Matsunaga K, Yamada A, Kikuchi M, Tsuda N, Ohta K, Takamori S, Yamana H, Fujita H, Itoh K: **Identification of a gene coding for a new squamous cell carcinoma antigen recognized by the CTL.** *J Immunol* 2000, **164**(5):2565–2574.

doi:10.1186/1743-422X-9-199

Cite this article as: Fukui *et al.*: Identification of B cell epitopes reactive to human papillomavirus type-16L1- derived peptides. *Virology Journal* 2012 **9**:199.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

