

RESEARCH

Open Access



# Function and diversity of P0 proteins among cotton leafroll dwarf virus isolates

Renan S. Cascardo<sup>1</sup>, Ighor L. G. Arantes<sup>1</sup>, Tatiane F. Silva<sup>2,3</sup>, Gilberto Sachetto-Martins<sup>1</sup>, Maité F. S. Vaslin<sup>2</sup> and Régis L. Corrêa<sup>1\*</sup>

## Abstract

**Background:** The RNA silencing pathway is an important anti-viral defense mechanism in plants. As a counter defense, some members of the viral family *Luteoviridae* are able to evade host immunity by encoding the P0 RNA silencing suppressor protein. Here we explored the functional diversity of P0 proteins among eight cotton leafroll dwarf virus (CLRVDV) isolates, a virus associated with a worldwide cotton disease known as cotton blue disease (CBD).

**Methods:** CLRVDV-infected cotton plants of different varieties were collected from five growing fields in Brazil and their P0 sequences compared to three previously obtained isolates. P0's silencing suppression activities were scored based on transient expression experiments in *Nicotiana benthamiana* leaves.

**Results:** High sequence diversity was observed among CLRVDV P0 proteins, indicating that some isolates found in cotton varieties formerly resistant to CLRVDV should be regarded as new genotypes within the species. All tested proteins were able to suppress local and systemic silencing, but with significantly variable degrees. All P0 proteins were able to mediate the decay of ARGONAUTE proteins, a key component of the RNA silencing machinery.

**Conclusions:** The sequence diversity observed in CLRVDV P0s is also reflected in their silencing suppression capabilities. However, the strength of local and systemic silencing suppression was not correlated for some proteins.

## Background

Cotton leafroll dwarf virus (CLRVDV) is the causal agent of an economically important cotton (*Gossypium hirsutum*) disease called cotton blue disease (CBD) [1]. *Aphis gossypii*-transmitted CBD has been observed in several cotton-producing areas of Central Africa, Asia and South America [2]. CBD symptoms are characterized by stunting, leaf rolling, vein yellowing, dark-green leaves and small bolls, leading therefore to severe yield losses when aphid populations are not properly controlled. In Brazil, CBD is present in almost all cotton growing fields and the disease was also partially controlled by the application of insecticides to decrease aphid populations and by the use of CBD-resistant cotton cultivars. Since 2006, several resistance breaking CLRVDV isolates have been observed throughout the country, producing CBD-like symptoms in formerly resistance cotton lines [3]. Apart from typical CBD symptoms, resistant or susceptible cotton varieties

infected with CLRVDV resistance-breaking isolates may also display reddish and withered leaves. Resistance breaking isolates are now widely distributed in Brazilian cotton growing areas, making the use of insecticide for aphid control compulsory.

The CLRVDV genome resembles a typical member of the genus *Polevirus*, family *Luteoviridae* and contains six open reading frames (ORF0 to ORF5) [4]. The genome is divided into two gene-containing portions, separated by an approximately 200 nucleotides intergenic region. Three open reading frames (ORF3, ORF4 and ORF5) are located in the 3'-end portion of the genome encoding for the structural proteins (capsid, movement and aphid-transmission proteins, respectively), while the 5'-end region of the genome encodes replication-related proteins (ORF1 and ORF2) and also a gene (ORF0) encoding the RNA silencing suppression protein P0. In general, the genome sequences of resistance breaking CLRVDV isolates are very similar to CLRVDV isolates from susceptible plants [3]. For example, the degree of sequence identity in all proteins encoded by ORFs 1 to 5 is greater than 93 % between two resistance breaking

\* Correspondence: regiscorrea@ufrj.br

<sup>1</sup>Department of Genetics, Federal University of Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil

Full list of author information is available at the end of the article

CLRVDV isolates (Ima2 and Acr3) and two non-resistance breaking ones (PV1 and ARG) in cotton plants. However, when the identities among P0 proteins are compared, the diversity is consistently higher, with identity numbers ranging from 85.8 to 86.6 % among the four isolates [3].

The P0 protein from several members of the genera *Polerovirus* and *Enamovirus*, family *Luteoviridae*, are known to be involved in the suppression of plant's anti-viral defense mechanisms at variable degrees, depending on the species and isolates [5–11]. P0's silencing suppression activity is mediated by promoting the destabilization of ARGONAUTE (AGO) proteins, key players in RNA silencing mechanisms [8, 12–15]. In plants, the RNA silencing pathway is triggered by double stranded RNAs (dsRNAs), which are processed by Dicer-like enzymes into small RNAs ranging from 20 to 24 nucleotides [16]. Viral-derived small interfering RNAs (siRNAs) produced during infections are readily recruited by AGO-containing RNA-induced silencing complexes (RISC) and used by the machinery to degrade viral genomic and sub-genomic sequences, being therefore an efficient anti-viral defense mechanism [17]. AGO is an important component of the machinery, since it directly binds to siRNAs and guide RISC to target RNAs. Viral RNA degradation may take place either at locally infected cells or at distal tissues, by the systemic movement of silencing signals [18]. A plethora of evolutionary unrelated viral proteins has evolved to cope with the anti-viral RNA silencing process. The P19 proteins from tombusviruses are one of the best characterized suppressors. P19 proteins are able to bind to siRNAs, preventing their loading into RISC [19]. Similarly, by degrading AGO proteins, P0 proteins are able to suppress the plant's anti-viral defense, allowing the infection to proceed. P0 proteins probably exert their activity through an F-box-dependent interaction with homologs of the S-phase kinase-related protein 1 (SKP1) ASK1 and ASK2 [20]. SKP1 is a core component of the SKP1/Cullin1/F-box (SCF) family of E3 ubiquitin ligases that mediate the ubiquitination of diverse regulatory and signaling proteins [21]. Point mutations in P0's F-box motif may abolish its interaction with SKP1 and consequentially decreasing AGO destabilization and viral pathogenesis [8, 9, 11, 20]. However, P0's activity is insensitive to proteasome inhibitors and the viral protein probably operates by hitchhiking cellular autophagy pathways endogenously used to modulate AGO homeostasis [13–15]. This model is supported by the increased accumulation of AGO proteins in the presence of autophagy inhibitors and by its co-localization with autophagic vesicles [14].

Recently, the P0 protein from an Argentinian isolate of CLRVDV (P0<sup>CL-ARG</sup>) has been characterized as a RNA silencing suppression protein [10]. The level of both local and systemic silencing suppression observed in P0<sup>CL-ARG</sup>


seems to be low when compared to other members of the group. Almost no suppression of systemic silencing is observed for P0<sup>CL-ARG</sup>, in line with what has been previously found for the P0 proteins from other members of the family [5, 10, 22]. However, it's known that P0 silencing suppression activity can vary even among closely related viruses. For example, European isolates of beet mild yellowing virus vary greatly in their ability to suppress local silencing [7]. Here, the local and systemic silencing suppression activities of P0 proteins from CLRVDV isolates collected in different parts of Brazil, including CBD resistant and susceptible cotton varieties, were assessed and compared to P0<sup>CL-ARG</sup>. Results indicate that silencing suppression capabilities are strain-specific and that strength of local and systemic silencing suppression is not correlated in CLRVDV P0 proteins.

## Results and discussion

### Sequence diversity among CLRVDV P0 proteins

When the genomic sequences of four previously identified CLRVDV isolates were compared, significant differences were only observed in the P0 coding sequence [3]. In order to better characterize the sequence diversity of P0 proteins among CLRVDV isolates, cotton plants from different varieties, displaying typical or atypical CBD symptoms were collected from different parts of Brazil. Sampled areas covered the main cotton producing areas of the country, most of them with significant geographical distance from each other (Fig. 1). In total, seven Brazilian CLRVDV isolates (PV1, Ima2, Acr9, Ipa4, Pm1, Hol1 and Pal3) were analyzed and compared to the Argentinian isolate (ARG) [4] and also to the P0 from an Australian isolate of potato leafroll virus (P0<sup>PL-AU</sup>) [8] (Table 1). The genomes from two of the Brazilian isolates analyzed (PV1 and Ima2) have been previously obtained [3, 23]. Three CLRVDV isolates (Acr9, Ima2 and Ipa4) were obtained from known cotton CLRVDV-resistant varieties and are therefore treated as resistance-breaking isolates. Two isolates (Ima2 and Pm1) were obtained from plants showing atypical symptoms (Table 1).

The amino acid sequence identity among CLRVDV P0s varied from 85.5 % (P0<sup>CL-Ima2</sup>/P0<sup>CL-Hol1</sup>) to 98.88 % (P0<sup>CL-Pm1</sup>/P0<sup>CL-Pal3</sup> and P0<sup>CL-ARG</sup>/P0<sup>CL-Pal3</sup>) (Table 2). The P0 sequences from the isolates Hol1, Pal3 and Pm1 are very close to the P0s from PV1 and ARG, the two isolates initially associated with CBD [1, 4], with identities ranging from 97.76 % to 98.88 % (Table 2). The isolates Acr9, Ima2 and Ipa4 also have a high sequence identity among them, ranging from 95.53 % to 97.2 %. The identity among P0 sequences from isolates Acr9, Ima2 and Ipa4, however, is lower than 90 % when compared to PV1 or ARG. The current taxonomic criteria in the family *Luteoviridae* states that viruses having amino acid divergence higher than 10 % in any protein



sequence should be considered as different species [24]. In this line, the isolates Acr9, Ima2 and Ipa4 should be regarded as a new species associated with CBD. Since P0 sequences are the most variable sequences among poleroviruses, it has been recently proposed that viruses having high diversity in this region, but with amino acid identities higher than 90 % in all other proteins should be regarded as genotypes of the same species and not as a different one [25, 26]. This kind of analysis can only be

made for the Ima2 isolate, the only one of the three with the genome fully sequenced [3]. But since the P0 sequences of the three isolates are very similar (Table 2), Acr9 and Ipa4 are probably also new genotypes of CLRDRV, as previously stated for Ima2 [3].

The separation of the isolates in two groups is also supported by phylogenetic analysis. It has been shown that, when full genomic sequences are not available, P0-based phylogenies can correctly reconstruct the relatedness

**Table 1** Brazilian isolates of cotton leafroll dwarf virus used in the study

Isolate	Location <sup>a</sup>	<i>G. hirsutum</i> cultivar	CBD resistance phenotype	Symptoms observed	Year
PV1	Primavera do Leste – MT	FM966	Susceptible	Typical	2004
Acr9	Acreuna – GO	CD406	Resistant	Typical	2006
Hol1	Holambra – SP	Nd	Nd	Typical	2007
Ima2	Campo Verde - MT	IAC25 RMD	Resistant	Atypical	2009
Ipa4	Ipameri - GO	Delta Opal	Resistant	Typical	2006
Pal3	Palotina - PR	CD034928	Nd	Typical	2006
Pm1	Patos de Minas - MG	Epamig1	Nd	Atypical	2007

<sup>a</sup>MT, Mato Grosso State; GO – Goiás State; SP – São Paulo State; PR – Paraná State; MG – Minas Gerais State

**Table 2** Percentage of amino acid identity among the viral isolates used in the study

P0 <sup>CL-Pal3</sup>	X									
P0 <sup>CL-Pm1</sup>	98.88	X								
P0 <sup>CL-PV1</sup>	98.51	98.14	X							
P0 <sup>CL-Hol1</sup>	98.51	98.14	97.76	X						
P0 <sup>CL-ARG</sup>	98.88	98.51	98.14	98.14	X					
P0 <sup>CL-Ima2</sup>	86.24	85.87	86.61	85.5	86.24	X				
P0 <sup>CL-Ipa4</sup>	88.1	87.73	87.73	87.36	87.36	96.28	X			
P0 <sup>CL-Acr9</sup>	88.1	87.73	87.73	87.36	88.1	95.53	97.02	X		
P0 <sup>PL-AU</sup>	18.21	18.21	17.84	18.21	18.58	17.84	18.58	18.58	X	
	P0 <sup>CL-Pal3</sup>	P0 <sup>CL-Pm1</sup>	P0 <sup>CL-PV1</sup>	P0 <sup>CL-Hol1</sup>	P0 <sup>CL-ARG</sup>	P0 <sup>CL-Ima2</sup>	P0 <sup>CL-Ipa4</sup>	P0 <sup>CL-Acr9</sup>	P0 <sup>PL-AU</sup>	


among poleroviruses [25]. A Neighbor-joining tree based on CLRDRV P0s groups the isolates Hol1, Pal3 and Pm1 with the two CBD founding members PV1 and ARG, while the isolates Acr9, Ima2 and Ipa4 clearly branch out, forming a well-supported group (Fig. 2). Since the three divergent isolates were all obtained from formerly cotton resistant varieties (Table 1), the silencing suppression activities from all isolates were then scored and compared.


**Suppression of local silencing by CLRDRV P0s**

Previous data have shown that P0<sup>CL-ARG</sup> is weak suppressor of local silencing when expressed in *Nicotiana benthamiana* leaves [10]. The suppression assay is based on the *Agrobacterium*-mediated transient co-expression of mGFP5 and a candidate silencing suppression protein in mGFP5-expressing *N. benthamiana* leaves (transgenic line 16c) [27]. When no silencing suppression is observed, the transiently expressed GFP triggers a strong RNA silencing response that ultimately leads to RNA degradation from both stable and transient transgenes. However, in the presence of a silencing suppression protein, GFP degradation is prevented and both transgenes (stable and transient) are expressed, increasing total GFP levels. The silencing suppression assay can be easily

monitored by hand UV lamps. In order to check whether the sequence diversity observed among the P0s could also reflect variable silencing suppression activities, the P0 from all seven Brazilian isolates and the Argentinian isolate were tested side-by-side in the 16c *N. benthamiana* line and compared to P0<sup>PL-AU</sup> and P19, two known suppressors of local silencing. All genes were cloned into the pGWB417 binary vector [28], leading to a 35S-driven expression of Myc-tagged proteins.

As expected, a red patch in the infiltrated area was observed when GFP was expressed in the absence of a silencing suppression protein (Fig. 3). Conversely, strong GFP fluorescence was observed in the presence of P19 or P0<sup>PL-AU</sup> in all time-points analyzed (Fig. 3). All tested CLRDRV P0s displayed obvious RNA silencing suppression activities. When scored based on GFP fluorescence, the levels of silencing suppression activity observed for the CLRDRV P0s were similar to P0<sup>PL-AU</sup>, but significantly lower than the P19 suppressor protein even at 3 days post-infiltration (dpi) (Fig. 3). The observed GFP fluorescence at 3 dpi correlates well with the accumulation of GFP RNAs when checked by real-time PCR at this time-point in *N. benthamiana* 16c plants (Fig. 4a). GFP RNA levels in GFP/P19-infiltrated plants were approximately 14 times higher than in control mock-infiltrated 16c plants. However, the GFP RNA fold change in P0<sup>PL-AU</sup> or CLRDRV P0-infiltrated plants, varied from only 2 to 6 times the levels obtained in the same control condition (Fig. 4a). At later time-points, GFP fluorescence started to fade at infiltrated areas of P0<sup>CL-Acr9</sup>, P0<sup>CL-Hol1</sup>, P0<sup>CL-Ima2</sup> and P0<sup>CL-Ipa4</sup>, indicating that those proteins are weaker suppressors than the other P0s (Fig. 3). The accumulation of the Myc-tagged suppressor proteins in the three biological replicates used for real-time PCR was checked by western blot using anti-Myc antibodies (Fig. 4b). Although all suppressors were expressed from the same vector background (pGWB417) [28], the P19 protein accumulated at levels consistently higher than the P0s. The strong suppression activity observed for P19 in the assay, therefore, might be correlated with its higher stability in *N.*





**Fig. 3** Suppression of local silencing by cotton leafroll dwarf virus P0s in *N. benthamiana* 16c plants. Transgenic *N. benthamiana* plants expressing GFP (line 16c) were co-infiltrated with *Agrobacterium* carrying plasmids to express GFP (silencing trigger) and candidate suppressor proteins (P0<sup>CL-PV1</sup>, P0<sup>CL-Acr9</sup>, P0<sup>CL-Hol1</sup>, P0<sup>CL-Ima2</sup>, P0<sup>CL-Ipa4</sup>, P0<sup>CL-Pal3</sup> and P0<sup>CL-Pm1</sup>). GFP co-expressed with empty vector was used as negative control. P0<sup>CL-ARG</sup>, P0<sup>PL-AU</sup> and P19 were used as positive controls in the assay. Pictures were taken at 3, 6 and 9 days post-infiltration (dpi)


*benthamiana* leaves. Since the P0s accumulated at similar levels, the observed differences in silencing suppression activities for those proteins might be due to functional divergence and not expression levels.

The fading phenotypes observed in 16c plants for the suppressor proteins P0<sup>CL-Acr9</sup>, P0<sup>CL-Hol1</sup>, P0<sup>CL-Ima2</sup> and P0<sup>CL-Ipa4</sup> were reproduced when the experiment was repeated in wild type plants (Additional file 1: Figure S1). When transiently expressed alone in wild type plants, the accumulation of GFP was lower than in the presence of the control strong suppressor P19, even at 3 dpi (Additional file 1: Figure S1). At 6 dpi, GFP was almost totally silenced when expressed alone, contrasting to what was observed in the presence of control constructs (P19 or PL<sup>P0-AU</sup>) or any of the CLRDRV P0s. From 6 dpi onwards, as mentioned before, GFP levels started to fade in the presence of P0<sup>CL-Acr9</sup>, P0<sup>CL-Hol1</sup>, P0<sup>CL-Ima2</sup> and P0<sup>CL-Ipa4</sup>, also indicating that those proteins are not able to suppress GFP silencing in *N. benthamiana* leaves for long periods. In all time-points analyzed, GFP accumulated at levels expressively higher when co-infiltrated with P19 than in the presence of any other P0, indicating that even the ones able to maintain GFP suppression at later times post-infiltration (P0<sup>PL-AU</sup>, P0<sup>CL-ARG</sup>, P0<sup>CL-PV1</sup>, P0<sup>CL-Pal3</sup> and P0<sup>CL-Pm1</sup>) should be regarded as moderate suppressors compared to the control used in the assays.

It has been previously shown that P0s depend on the presence of a F-box-like motif to exert their silencing suppression activity [8, 9, 11, 20]. The hallmark amino acids LPxx(L/I)x<sup>10-13</sup>P could be found in all CLRDRV P0s tested (Additional file 2: Figure S2). However, isoleucine is changed to an amino acid with similar biochemical properties (valine) in the three resistance-breaking CLRDRV isolates (Acr9, Ima2 and Ipa4). The ring structure amino acids known to affect local silencing suppression activity of melon aphid-borne yellows virus P0 are also present and conserved among the CLRDRV P0s (Additional file 2: Figure S2) [9]. Therefore, the local silencing suppression variability observed among CLRDRV P0s is probably associated with alternative functional residues.

**Suppression of systemic silencing by CLRDRV P0s**

Viral siRNAs produced in infected cells may also move systemically through vascular tissues to reach other parts



of the plants [18]. In the *N. benthamiana* 16c assay, systemic silencing can be visualized by the appearance of red-silenced areas especially around the veins of newly developed leaves. Eventually, silencing signals may spread throughout the leaves, producing completely silenced plants. The systemic silencing suppression activities of all CLRDRV P0s were scored and compared to the strong systemic suppressor P0<sup>PL-AU</sup> [8]. At 16 dpi, 15 out of 20 plants assayed showed systemic silencing when infiltrated with the GFP silencing-trigger construct in the absence of any suppressor protein and almost 100 % of the plants were silenced by 20 dpi (Table 3). As expected, P0<sup>PL-AU</sup> completely blocked the spread of silencing signals as no silenced plants were observed at 16, 20 or 29 dpi. However, the suppression of systemic silencing mediated by the CLRDRV P0s varied among the different isolates. In our experimental conditions, only 3 plants out of 20 were silenced at 16 dpi when co-infiltrated with P0<sup>CL-ARG</sup> (Table 3). The number of silenced plants, however, increased to 7 and 10 out of 20 at 20 dpi and 29 dpi, respectively, indicating that P0<sup>CL-ARG</sup> is

a moderate suppressor of silencing signals (Table 3). This result contrast to what has been previously observed for P0<sup>CL-ARG</sup>, where 8 out 10 plants were already silenced by 15 dpi in the presence of the protein [10]. Since the spread of RNA silencing signals may be

**Table 3** Proportion of plants showing systemic silencing at 16, 20 and 29 days post-infiltration (dpi)

Infiltration	16 dpi	20 dpi	29 dpi
GFP + Vector	15/20	19/20	19/20
GFP + P0 <sup>PL-AU</sup>	0/20	0/20	0/20
GFP + P0 <sup>CL-PV1</sup>	0/20	0/20	0/20
GFP + P0 <sup>CL-ARG</sup>	3/20	7/20	10/20
GFP + P0 <sup>CL-Acr9</sup>	5/20	7/20	7/20
GFP + P0 <sup>CL-Hol1</sup>	0/20	1/20	2/20
GFP + P0 <sup>CL-Ima2</sup>	7/20	9/20	11/20
GFP + P0 <sup>CL-Ipa4</sup>	0/20	0/20	0/20
GFP + P0 <sup>CL-Pal3</sup>	0/20	0/20	0/20
GFP + P0 <sup>CL-Pm1</sup>	1/20	1/20	1/20

influenced by environmental conditions [29–31] and possibly by the number of infiltrated leaves, concentration and strain of *Agrobacterium* used in the assay and vector background, the difference in the systemic silencing activity observed for  $P0^{CL-ARG}$  might be due to different experimental settings.

In line with what has been observed for  $P0^{CL-ARG}$ , the  $P0$  proteins  $P0^{CL-Acr9}$  and  $P0^{CL-Ima2}$  were also moderate suppressors of systemic silencing (Table 3). However, the  $P0$  proteins from five isolates ( $PV1$ ,  $Hol1$ ,  $Ipa4$ ,  $Pal3$  and  $Pm1$ ) efficiently blocked the spread of systemic silencing signals, with a silencing suppression activity similar to  $P0^{PL-AU}$  (Table 3). During the experiments, 70 % of the plants infiltrated with suppressor proteins  $P0^{CL-Pal3}$  and  $P0^{CL-Pm1}$  and 50 % of  $P0^{PL-AU}$ -infiltrated ones displayed strong necrotic lesions at late times after infiltration, most of them starting at 10 dpi (data not shown). Therefore, the suppression of systemic silencing by those proteins could have also been influenced by the induced cell death. Similar necrotic phenotypes have also been observed for  $P0^{PL-AU}$  [8] and for the  $P0$ s from sugarcane yellow leaf virus and beet western yellows virus [6].


#### AGO destabilization by CLRDRV P0s

The  $P0$  proteins from some members of the family *Luteoviridae* are able to destabilize the expression of AGO proteins [8, 11–15]. However, the activity of CLRDRV  $P0$  in AGO decay has never been tested. For that, a Myc-tagged version of the *Arabidopsis thaliana* AGO1 protein, known to be involved in several RNA silencing pathways, including anti-viral defense [32, 33], was transiently expressed via *Agrobacterium* infiltration in wild type *N. benthamiana* leaves in the presence or absence of different  $P0$ s or control constructs. The accumulation of AtAGO1-Myc protein was detected by anti-myc antibodies when co-expressed without  $P0$  suppressors or in the presence of  $P19$  (Fig. 5). All  $P0$ s, including  $P0^{PL-AU}$ , the seven Brazilian

isolates and the Argentinian isolate of CLRDRV were able to strongly decrease the AtAGO1-Myc levels when co-expressed. Therefore, the local and systemic silencing differences observed among the isolates might be due to the regulation of still unknown cellular proteins by  $P0$ , possibly other AGO members [8], or due to small differences in AGO1 accumulation that are not detected due to western blot resolution limits.

#### Conclusions

Our results indicated a high diversity among  $P0$  proteins from Brazilian and Argentinian isolates of CLRDRV, a virus associated with CBD. All CLRDRV  $P0$  proteins analyzed were able to mediate AtAGO1 decay, however, variable silencing suppression activities were observed, probably reflecting their sequence diversity.  $P0^{CL-ARG}$  was a moderate silencing suppressor of both local and systemic silencing in our experiments, when compared to the positive control constructs used in the assays (Figs. 3, 4 and Additional file 1: Figure S1). Three proteins ( $P0^{CL-PV1}$ ,  $P0^{CL-Pal3}$  and  $P0^{CL-Pm1}$ ) were also moderate suppressors of local silencing, but strong suppressors of systemic silencing. Four other proteins behaved as weak suppressors of local silencing. Contrasting to control constructs ( $P19$  and  $P0^{PL-AU}$ ) and to other CLRDRV  $P0$ s, those four proteins ( $P0^{CL-Acr9}$ ,  $P0^{CL-Ima2}$ ,  $P0^{CL-Hol1}$  and  $P0^{CL-Ipa4}$ ) could not support GFP suppression for long periods when assayed in the mGFP5-expressing *N. benthamiana* 16c line (Fig. 3) or in wild type plants (Additional file 1: Figure S1). GFP levels clearly started to fade in the presence of those proteins from 6 dpi onwards (Fig. 3 and Additional file 1: Figure S1). However, two of the weak local silencing suppressor proteins ( $P0^{CL-Hol1}$  and  $P0^{CL-Ipa4}$ ) were able to almost completely block the spread of systemic silencing signals when assayed in 16c transgenic lines. Despite of their weak local silencing,  $P0^{CL-Hol1}$  and  $P0^{CL-Ipa4}$  are as strong as the control  $P0^{PL-AU}$



**Fig. 5** Accumulation of a Myc-tagged version of the *A. thaliana* AGO1 protein transiently expressed in *N. benthamiana* leaves in the presence or absence of  $P0$  proteins. Myc-tagged versions of cotton leafroll dwarf virus  $P0$ s ( $P0^{CL-ARG}$ ,  $P0^{CL-PV1}$ ,  $P0^{CL-Acr9}$ ,  $P0^{CL-Hol1}$ ,  $P0^{CL-Ima2}$ ,  $P0^{CL-Ipa4}$ ,  $P0^{CL-Pal3}$  or  $P0^{CL-Pm1}$ ) and potato leafroll virus  $P0$  ( $P0^{PL-AU}$ ) were used in the assay. Non-infiltrated plants (NI) were used as negative controls and AtAGO1 infiltrated with the empty vector or with a Myc-tagged  $P19$  were used as positive controls. Membranes were probed with anti-Myc antibodies. Ponceau staining was used as a loading control. All co-infiltrations were performed in the presence of  $P19$  (without tag) to stabilize mRNA *Agrobacterium*-mediated experiments

protein in suppressing systemic silencing. It's tempting to speculate therefore that the strength of local and systemic silencing suppression activity might be genetically unlinked in P0 proteins. Furthermore, these data indicate that the silencing suppression capabilities of the distinct CLRDV P0 proteins are not directly linked to their genetic diversity.

## Methods

### Plant material and DNA constructs

*Gossypium hirsutum* plants belonging to at least six cultivars (FM966, CD406, CD034928, IAC25 RMD, Delta Opal and Epamig1) were collected in five different States of Brazil (Goiás, Mato Grosso, Minas Gerais, Paraná and São Paulo) (Table 1 and Fig. 1). The Ima2 isolate was collected in Campo Verde – Mato Grosso, but passed through the IAC24 RMD cotton variety at Instituto Matogrossense do Algodão (Primavera do Leste – Mato Grosso) before being sent for analysis [34]. Harvesting and maintenance of plants were performed according to Brazilian rules (MP 2.186-16/2001). Total RNA of all plants were extracted using Qiagen Plant RNA kit and 2.5 µg were used to prepare cDNAs with the O5R2 primer (5'-GCAACCTTTTATAGTCTCTCCAAT-3'), which anneals in the middle of CLRDV ORF5. The ORF0 sequences from all Brazilian isolates were amplified with primers CLP0\_F (5'-CACCATGTTGAATTTGATCATC TGCAG-3') and CLP0\_R (5'-ACTGCTTTCTCCTTCAC-3') and cloned into pENTRY-D-TOPO (Invitrogen). The ORF0 of the Argentinian isolate of CLRDV [GenBank: NC\_014545.1] was synthesized and cloned into a pUC plasmid by the Blue Heron Biotechnology Inc (USA). The Argentinian ORF0 [10] and the P19 coding sequence [35] were amplified with primers CLP0\_TOPO\_F/CLP0\_R and P19\_TP\_F (5'-CACCATGGAACGAGCTATAACAAGGAA ACG -3')/P19\_R (5'-TTACTCGCTTTCTTTTTCGAAG G-3'), respectively, and also cloned into pENTRY-D-TOPO (Invitrogen). All amplifications were performed with the Phusion High Fidelity Polymerase (NEB). Entry vectors containing the *Arabidopsis thaliana* Ago1 coding sequence [TAIR: AT1G48410] and the P0 from the Australian isolate of PLRV [GenBank: D13953.1] were described previously [8].

Genes in entry gateway clones were sequenced in both directions in automated ABI sequencers through dye terminator cycle method, using primers annealing in vector sequences. The accession numbers for the new P0 sequences obtained here are: [GenBank:KR185733] (isolate Acr9), [GenBank:KR185734] (isolate Pm1), [GenBank:KR185735] (isolate Hol1), [GenBank:KR185736] (isolate Ipa4), [GenBank:KR185737] (isolate Pal3). All genes in entry vectors were transferred through LR reactions to the binary destination vector pGWB417 [28],

resulting in 35S-driven, Myc-tagged proteins when expressed in plants.

### Sequence analysis

Multiple sequence alignments of deduced amino acid sequences were performed with ClustalW2 (<http://www.ebi.ac.uk/Tools/msa/clustalw2/>) and phylogenetic reconstructions were performed with the MEGA 4 software [36]. Trees were constructed by the neighbor-joining (NJ) method [37], with the pair-wise deletion option and number of differences matrix.

### Agroinfiltration

*Agrobacterium tumefaciens*, strain GV3101, were infiltrated in *Nicotiana benthamiana* leaves as described previously [38]. Cells were individually diluted to an optical density of 1.0 at 600 nm before mixing the cultures. Leaves were infiltrated in the abaxial surfaces with needleless syringes and the infiltrated plants were incubated in growth chambers with a 16-hour photoperiod at 24 °C.

For the local silencing suppression assay, three leaves of 5-weeks-old mGFP5-expressing *N. benthamiana* plants (wild type or 16c line) [27] were co-infiltrated with equal volumes of *A. tumefaciens* harboring plasmids expressing mGFP5 and pGWB417 with or without candidate silencing suppressor genes. For the systemic silencing suppression assay, only one leaf of 3-weeks-old *N. benthamiana* 16c plants were co-infiltrated with equal volumes of *A. tumefaciens* expressing mGFP5 and pGWB417 (negative control) or with pGWB417 expressing candidate silencing suppressor genes. GFP fluorescence was observed under a long-wavelength UV lamp and the number of plants having systemic silencing scored in different time-points.

For the AGO1 destabilization assay, three leaves of 5-week-old wild type *N. benthamiana* plants were infiltrated with *A. tumefaciens* harboring plasmids pJL3:P19 [35], pGWB417-AtAGO1-MYC, and pGWB417 with or without candidate silencing suppression genes in a proportion of 30 %, 35 % and 35 %, respectively. The infiltrated leaves were collected 4 days after infiltration.

### qRT-PCR

Total RNA from approximately 100 mg of infiltrated *N. benthamiana* leaf tissue was extracted using the Plant RNA Purification Reagent (Invitrogen) according to manufacturer's instructions. The quality of the RNA was checked by electrophoresis on 1.0 % agarose gels in 0.5X TAE buffer, and the RNA was quantified with NanoDrop 2000 spectrophotometer (Thermo Scientific). One microgram of total RNA was treated with DNase I (Promega) according to manufacturer's instructions and used for cDNA synthesis with oligo d(T) primer and the SuperScriptIII (Invitrogen) enzyme, according to manufacturer's

instructions. Quantitative PCR reactions were performed in a total volume of 20  $\mu$ L, using 5  $\mu$ L of a 20-fold diluted cDNA. The amplification reactions were performed using the SYBR<sup>®</sup> Select Master Mix (Applied Biosystems), according to manufacturer's instructions. Primers used for GFP were qmGFP5\_F3 (5'-AGTGGAGAGGGTGAAGGTGATGC-3') and qmGFP5\_R4 (5'-TCCCTCAGGCATGGCGCTCTT-3'). The genes Ubiquitin3 (Ubi3) and Elongation factor-1  $\alpha$  (EF-1) were used as reference genes, with primers previously described [39]. Three biological and technical replicates were used for all samples. Quantification of GFP expression levels was performed using the comparative CT method ( $\Delta\Delta$ CT) through the Miner and qBase softwares [40–42]. The t-student test was performed to compare the samples.

### Western blotting

Infiltrated *Nicotiana benthamiana* leaves were ground in liquid nitrogen and mixed with sample buffer (100 mM Tris [pH 6.8], 20 % glycerol, 4 % SDS, and 0.2 % bromophenol blue) containing 10 %  $\beta$ -mercaptoethanol [43]. Samples were then boiled at 90 °C for 10 min, and centrifuged for 5 min at 13,000  $\times$  g before loading on a gel. Extracts were run in 8 % SDS-PAGE gels for the detection of AtAGO1-Myc and in 12 % SDS-PAGE gels for detecting P19-myc, P0<sup>PL-AU</sup>-myc and P0<sup>CLS</sup>-myc with anti-myc antibody (1:2,000; Sigma, clone 9E10), followed by an anti-mouse HRP secondary antibody (1:5,000; Bio-Rad). Antibody–protein interactions were visualized using an enhanced chemiluminescence detection kit (GE Healthcare) according to the manufacturer's instructions.

### Additional files

**Additional file 1: Figure S1.** Suppression of local silencing by cotton leafroll dwarf virus P0s in *N. benthamiana* wild type plants. *N. benthamiana* plants were co-infiltrated with *Agrobacterium* carrying plasmids to express GFP and candidate suppressor proteins (P0<sup>CL-PV1</sup>, P0<sup>CL-Acr9</sup>, P0<sup>CL-Hol1</sup>, P0<sup>CL-lma2</sup>, P0<sup>CL-tpa4</sup>, P0<sup>CL-Pal3</sup> and P0<sup>CL-Pm1</sup>). GFP co-expressed with empty vector was used as negative control. P0<sup>CL-ARG</sup>, P0<sup>PL-AU</sup> and P19 were used as positive controls in the assay. Pictures were taken at 3, 6, 9 and 12 days post-infiltration (dpi). (PNG 4388 kb)

**Additional file 2: Figure S2.** Amino acid sequence alignment of cotton leafroll dwarf virus P0s. Sequences were aligned with ClustalW2 and conserved amino acids were shaded in black with Mview. The red bar highlights the conserved F-box-like domain. Functional ring structure residues described in Han et al., 2010 are marked with asterisks. (PNG 1391 kb)

### Competing interests

The author(s) declare that they have no competing interests.

### Authors' contributions

RSC, GSM, MFSV and RLC conceived and designed the experiments. RSC, ILGA and TFS performed the experiments. MFSV and TFS contributed with plant material. GSM, MFSV and RLC contributed reagents/materials/analysis tools. RSC and RLC wrote the manuscript. All authors read and approved the final manuscript.

### Acknowledgements

Authors acknowledge Dr. Jean-Lois Bélot and Dr. Rafael Galbieri, from Instituto Matogrossense do Algodão and Dr. Nelson Suassuna from EMBRAPA Algodão for sending the cotton infected plants. This study is part of the thesis research of RSC in pursuit of his PhD in Genetics at the Genetics Department of the Federal University of Rio de Janeiro. This work was supported by grants from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) to GSM, MFSV and RLC. RSC and ILGA were supported by fellowships from CNPq.

### Author details

<sup>1</sup>Department of Genetics, Federal University of Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil. <sup>2</sup>Department of Virology, Federal University of Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil. <sup>3</sup>Present address: Departamento de Biotecnologia, Escola de Engenharia de Lorena, Universidade de São Paulo, Lorena, São Paulo, Brazil.

Received: 28 April 2015 Accepted: 4 August 2015

Published online: 12 August 2015

### References

- Correa RL, Silva TF, Simoes-Araujo JL, Barroso PA, Vidal MS, Vaslin MF. Molecular characterization of a virus from the family Luteoviridae associated with cotton blue disease. *Arch Virol.* 2005;150:1357–67.
- Cauquil J. Etudes sur une maladie d'origine virale du cotonnier: la maladie bleue. *Cot Fibres Trop.* 1977;32:259–78.
- Da Silva AKF, Romanel E, Silva T da F, Castilhos Y, Schrago CG, Galbieri R, et al. Complete genome sequences of two new virus isolates associated with cotton blue disease resistance breaking in Brazil. *Arch Virol.* 2015;160:1371–4.
- Distéfano AJ, Kresic IB, Hopp HE. The complete genome sequence of a virus associated with cotton blue disease, cotton leafroll dwarf virus, confirms that it is a new member of the genus Ploverovirus. *Arch Virol.* 2010;155:1849–54.
- Pfeffer S, Dunoyer P, Heim F, Richards KE, Jonard G, Ziegler-Graff V. P0 of beet Western yellows virus is a suppressor of posttranscriptional gene silencing. *J Virol.* 2002;76:6815–24.
- Mangwende T, Wang ML, Borth W, Hu J, Moore PH, Mirkov TE, et al. The P0 gene of Sugarcane yellow leaf virus encodes an RNA silencing suppressor with unique activities. *Virology.* 2009;384:38–50.
- Kozłowska-Makulska A, Guillely H, Szyndel MS, Beuve M, Lemaire O, Herrbach E, et al. P0 proteins of European beet-infecting poleroviruses display variable RNA silencing suppression activity. *J Gen Virol.* 2010;91(Pt 4):1082–91.
- Fusaro AF, Correa RL, Nakasugi K, Jackson C, Kawchuk L, Vaslin MFS, et al. The Enamovirus P0 protein is a silencing suppressor which inhibits local and systemic RNA silencing through AGO1 degradation. *Virology.* 2012;426:178–87.
- Han YH, Xiang HY, Wang Q, Li YY, Wu WQ, Han CG, et al. Ring structure amino acids affect the suppressor activity of melon aphid-borne yellows virus P0 protein. *Virology.* 2010;406:21–7.
- Delfosse VC, Agrofolgio YC, Casse MF, Kresic IB, Hopp HE, Ziegler-Graff V, et al. The P0 protein encoded by cotton leafroll dwarf virus (CLRVDV) inhibits local but not systemic RNA silencing. *Virus Res.* 2014;180:70–5.
- Zhuo T, Li YY, Xiang HY, Wu ZY, Wang XB, Wang Y, et al. Amino acid sequence motifs essential for P0-mediated suppression of RNA silencing in an isolate of potato leafroll virus from Inner Mongolia. *Mol Plant Microbe Interact.* 2014;27:515–27.
- Bortolamiol D, Pazhouhandeh M, Marrocco K, Genschik P, Ziegler-Graff V. The Ploverovirus F box protein P0 targets ARGONAUTE1 to suppress RNA silencing. *Curr Biol.* 2007;17:1615–21.
- Baumberger N, Tsai CH, Lie M, Havecker E, Baulcombe DC. The Ploverovirus silencing suppressor P0 targets ARGONAUTE proteins for degradation. *Curr Biol.* 2007;17:1609–14.
- Derrien B, Baumberger N, Schepetilnikov M, Viotti C, De Cillia J, Ziegler-Graff V, et al. Degradation of the antiviral component ARGONAUTE1 by the autophagy pathway. *Proc Natl Acad Sci U S A.* 2012;109:15942–6.
- Csorba T, Loza R, Hutvagner G, Burgyan J. Ploverovirus protein P0 prevents the assembly of small RNA-containing RISC complexes and leads to degradation of ARGONAUTE1. *Plant J.* 2010;62:463–72.

16. Bologna NG, Voinnet O. The diversity, biogenesis, and activities of endogenous silencing small RNAs in Arabidopsis. *Annu Rev Plant Biol.* 2014;65:473–503.
17. Incarbone M, Dunoyer P. RNA silencing and its suppression: Novel insights from in planta analyses. *Trends Plant Sci.* 2013;18:382–92.
18. Melnyk CW, Molnar A, Baulcombe DC. Intercellular and systemic movement of RNA silencing signals. *EMBO J.* 2011;30:3553–63.
19. Lakatos L, Csorba T, Pantaleo V, Chapman EJ, Carrington JC, Liu YP, et al. Small RNA binding is a common strategy to suppress RNA silencing by several viral suppressors. *EMBO J.* 2006;25:2768–80.
20. Pazhouhandeh M, Dieterle M, Marrocco K, Lechner E, Berry B, Brault V, et al. F-box-like domain in the polerovirus protein P0 is required for silencing suppressor function. *Proc Natl Acad Sci.* 2006;103:1994–9.
21. Hua Z, Vierstra RD. The cullin-RING ubiquitin-protein ligases. *Annu Rev Plant Biol.* 2011;62:299–334.
22. Guilley H, Bortolamiol D, Jonard G, Bouzoubaa S, Ziegler-Graff V. Rapid screening of RNA silencing suppressors by using a recombinant virus derived from Beet necrotic yellow vein virus. *J Gen Virol.* 2009;90:2536–41.
23. De Andrade RRS, Vaslin MFS. SearchSmallRNA: a graphical interface tool for the assemblage of viral genomes using small RNA libraries data. *Viro J.* 2014;11:45.
24. Domier LL. *Virus Taxonomy, Ninth Report of the International Committee on Taxonomy of Viruses.* London: Elsevier/Academic Press; 2012.
25. Lin YH, Gao SJ, Damaj MB, Fu HY, Chen RK, Mirkov TE. Genome characterization of sugarcane yellow leaf virus from China reveals a novel recombinant genotype. *Arch Virol.* 2014;159:1421–9.
26. Knierim D, Tsai WS, Maiss E, Kenyon L. Molecular diversity of poleroviruses infecting cucurbit crops in four countries reveals the presence of members of six distinct species. *Arch Virol.* 2014;159:1459–65.
27. Brigneti G, Voinnet O, Li WX, Ji LH, Ding SW, Baulcombe DC. Viral pathogenicity determinants are suppressors of transgene silencing in *Nicotiana benthamiana*. *EMBO J.* 1998;17:6739–46.
28. Nakagawa T, Suzuki T, Murata S, Nakamura S, Hino T, Maeo K, et al. Improved Gateway binary vectors: high-performance vectors for creation of fusion constructs in transgenic analysis of plants. *Biosci Biotechnol Biochem.* 2007;71:2095–100.
29. Szittyá G, Silhavy D, Molnar A, Havelda Z, Lovas A, Lakatos L, et al. Low temperature inhibits RNA silencing-mediated defence by the control of siRNA generation. *EMBO J.* 2003;22:633–40.
30. Qu F, Ye X, Hou G, Sato S, Clemente TE, Morris TJ. RDR6 has a broad-spectrum but temperature-dependent antiviral defense role in *Nicotiana benthamiana*. *J Virol.* 2005;79:15209–17.
31. Kotakis C, Vrettos N, Kotsis D, Tsagris M, Kotzabasis K, Kalantidis K. Light intensity affects RNA silencing of a transgene in *Nicotiana benthamiana* plants. *BMC Plant Biol.* 2010;10:220.
32. Qu F, Ye X, Morris TJ. Arabidopsis DRB4, AGO1, AGO7, and RDR6 participate in a DCL4-initiated antiviral RNA silencing pathway negatively regulated by DCL1. *Proc Natl Acad Sci U S A.* 2008;105:14732–7.
33. Baumberger N, Baulcombe DC. Arabidopsis ARGONAUTE1 is an RNA Slicer that selectively recruits microRNAs and short interfering RNAs. *Proc Natl Acad Sci.* 2005;102:11928–33.
34. Galbieri R, Cia E, Fuzatto M, Franzon R, Belot J, Dias JC. Transmissibility and cotton genotypes' response to an atypical form of vein mosaic virus disease. *Trop Plant Pathol.* 2010;35:88–95.
35. Lindbo JA. High-efficiency protein expression in plants from agroinfection-compatible Tobacco mosaic virus expression vectors. *BMC Biotechnol.* 2007;7:52.
36. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol.* 2007;24:1596–9.
37. Saitou N, Saitou N, Nei M, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol.* 1987;4:406–25.
38. English JJ, Davenport GF, Elmayer T, Vaucheret H, Baulcombe DC. Requirement of sense transcription for homology-dependent virus resistance and trans-inactivation. *Plant J.* 1997;12:597–603.
39. Rotenberg D, Thompson TS, German TL, Willis DK. Methods for effective real-time RT-PCR analysis of virus-induced gene silencing. *J Virol Methods.* 2006;138:49–59.
40. Livak KJ, Livak KJ, Schmittgen TD, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta Ct}$  Method. *Methods.* 2001;25:402–8.
41. Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J. qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. *Genome Biol.* 2007;8:R19.
42. Zhao S, Fernald RD. Comprehensive algorithm for quantitative real-time polymerase chain reaction. *J Comput Biol.* 2005;12:1047–64.
43. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 1970;227:680–5.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

