

RESEARCH

Open Access



# Avian influenza H9N2 virus isolated from air samples in LPMs in Jiangxi, China

Xiaoxu Zeng<sup>1†</sup>, Mingbin Liu<sup>2†</sup>, Heng Zhang<sup>1</sup>, Jingwen Wu<sup>2</sup>, Xiang Zhao<sup>1</sup>, Wenbing Chen<sup>1</sup>, Lei Yang<sup>1</sup>, Fenglan He<sup>2</sup>, Guoyin Fan<sup>2</sup>, Dayan Wang<sup>1</sup>, Haiying Chen<sup>2\*</sup> and Yuelong Shu<sup>1\*</sup>

## Abstract

**Background:** Recently, avian influenza virus has caused repeated worldwide outbreaks in humans. Live Poultry Markets (LPMs) play an important role in the circulation and reassortment of novel Avian Influenza Virus (AIVs). Aerosol transmission is one of the most important pathways for influenza virus to spread among poultry, from poultry to mammals, and among mammals.

**Methods:** In this study, air samples were collected from LPMs in Nanchang city between April 2014 and March 2015 to investigate possible aerosol transmission of AIVs. Air samples were detected for Flu A by Real-Time Reverse Transcription-Polymerase Chain Reaction (RRT-PCR). If samples were positive for Flu A, they were inoculated into 9- to 10-day-old specific-pathogen-free embryonated eggs. If the result was positive, the whole genome of the virus was sequenced by MiSeq. Phylogenetic trees of all 8 segments were constructed using MEGA 6.05 software.

**Results:** To investigate the possible aerosol transmission of AIVs, 807 air samples were collected from LPMs in Nanchang city between April 2014 and March 2015. Based on RRT-PCR results, 275 samples (34.1%) were Flu A positive, and one virus was successfully isolated with embryonated eggs. The virus shared high nucleotide homology with H9N2 AIVs from South China.

**Conclusions:** Our study provides further evidence that the air in LPMs can be contaminated by influenza viruses and their nucleic acids, and this should be considered when choosing and evaluating disinfection strategies in LPMs, such as regular air disinfection. Aerosolized viruses such as the H9N2 virus detected in this study can increase the risk of human infection when people are exposed in LPMs.

**Keywords:** Influenza, Type a, H9N2, Aerosol, Phylogenetic, Transmission

## Background

While aquatic birds are the natural hosts of avian influenza virus (AIVs), the transmission of AIVs from wild birds to poultry has been seen in many places in the world, and the virus has caused frequent sporadic human outbreaks owing to contacts between poultry and humans [1–6]. During the past 20 years, live poultry markets (LPMs), which are widely distributed in China

and other developing countries, were proved to be the major source of human infections with AIVs, as well as a place for potential AIV reassortment and interspecies transmission [7, 8].

In previous studies searching for the source of human infection in LPMs, close attention was paid to poultry throat and cloaca swabs, as well as environmental samples such as feces, feathers, and poultry drinking water, while aerosol samples in LPMs were poorly investigated. Aerosol transmission is one of the most important pathways for influenza virus to spread among poultry, from poultry to mammals, and among mammals [9–11]. In recent years, human infection with many subtypes of AIVs has been reported, including H5N1, H5N6, H7N9, H10N8, H6N1, and H9N2 AIVs, and field epidemiological investigation showed some patients had no

\* Correspondence: nccdchyl@126.com; yshu@cnic.org.cn

†Equal contributors

<sup>2</sup>Nanchang Center for Disease Control and Prevention, Nanchang city 330038, People's Republic of China

<sup>1</sup>Chinese National Influenza Center, National Institute for Viral Disease Control and Prevention, Collaboration Innovation Center for Diagnosis and Treatment of Infectious Diseases, Chinese Center for Disease Control and Prevention, Key Laboratory for Medical Virology, National Health and Family Planning Commission, 155 Changbai Road, Beijing 102206, People's Republic of China


history of direct contact with any live poultry before illness onset [3, 12, 13]. For example, in late 2013, the first human infection with avian influenza H10N8 virus was reported in Nanchang city, Jiangxi province, China. The patient had visited the market 4 days before the onset of disease without touching any poultry [2]. To study possible aerosol transmission of AIVs in LPMs, air samples were collected from LPMs in Nanchang city between April 2014 and March 2015.


## Methods

### Sample collection

In this study, 46 sample sites in ten different LPMs were selected in Nanchang city, Jiangxi province, China. These LPMs were distributed in four districts (Xinjian, Donghu, Xihu, and Qingshanhu) (Fig. 1) of the city. Jingdong Bird and Flower market in Qingshanhu district was a large wholesale market; the others were all retail markets. Air samples were collected monthly from April 2014 to March 2015 via special air pumps (SQC-1000,

Yancheng Tianyue Instrument & Meter Co., Ltd., Jiangsu, China). This equipment is widely applied in collecting aerial particulates and aerosols in China and uses a collection bottle (Qingdao Junray Intelligent Instrument Co., Ltd.) to collect air samples by passing through a transport medium. In each month, 10 air samples were collected in Jingdong Bird and Flower market, and 6 air samples were collected in each of the other retail LPMs. For retail markets, first, the wind direction in the market was determined. The poultry stall was chosen as the study center; 2 air samples were collected on the left and right of the stall at 3 m from the stall; and three other air samples were collected at 3 m, 5 m and 7 m from the stall in the downwind direction; another air sample was collected in the upwind direction at 1 m distance from the stall (Fig. 2a). For Jingdong Bird and Flower wholesale market, five air samples were collected in the east, south, west, and north and the center of the LPM. Samples were also collected 1 m,





3 m, 5 m and 7 m from the outside entrance (Fig. 2b). Each air sample was collected for 1.5 h, then stored in MEM (Gibco), aliquoted into 3 tubes and stored at  $-80^{\circ}\text{C}$ . To avoid excess bubbles, no BSA was included in the transport medium during sample collection; instead, BSA was added immediately after collection. All samples were transported on ice.

#### Detection of influenza A nucleic acid and virus isolation

RNA was extracted with Qiagen RNeasy Mini kit, and influenza A was detected by RRT-PCR with primer and probe sets recommended by WHO (FluA-Forward: GAC CRATCCTGTCACCTCTGAC; FluA-Reverse: GGGCATT YTGACAAAKCGTCTACG;

FluA-probe: TGCAGTCCTCGCTCACTGGGCACG). If the aerosol samples were Flu A-positive, they were treated with 6 antibiotics (Penicillin:  $2 \times 10^6$  U/L, Streptomycin: 200 mg/L, Polymyxin:  $2 \times 10^6$  U/L, Gentamicin: 250 mg/L, Nystatin:  $0.5 \times 10^6$  U/L, Levofloxacin Hydrochloride: 600 mg/L) for 1 h, then virus isolation was conducted by inoculating the treated samples into 9- or 10-day-old specific-pathogen-free embryonated eggs. After incubation at  $37^{\circ}\text{C}$  for 48 h, the allantoic liquid of the eggs was harvested and tested by Flu A fast diagnostic kits (Colloidal Gold Antibody Conjugates, Beijing NBGen Bio-technology Limited Company). If the result was positive, the allantoic liquid was aliquoted and stored at  $-80^{\circ}\text{C}$ .

#### Whole-genome sequencing of influenza virus with MiSeq NGS platform

Viral RNA was extracted from harvested allantoic liquids using the Qiagen RNeasy Mini kit. Invitrogen SuperScript III One-step RT-PCR platinum Taq HiFi kit was used to amplify the RNA with one-step RT-PCR primers as follows: influenza primer A, GGGGGGAGCAAAAAGCAGG; influenza primer B, GGGGGGAGCGAAAAGCAGG influenza primer C, CGGGTTATTAGTAGAAACAAGG [14]. PCR products were purified by QIAGEN MinElute Reaction Cleanup Kit. One ng of DNA product was processed for NGS (next generation sequencing) sample preparation with the Illumina Nextera XT DNA Sample Preparation Kit (96 Samples), and sequencing was performed using a MiSeq v2 kit (500 cycles) to produce  $2 \times 250$  paired-end reads (Illumina, San Diego, CA, USA) [15]. After automated cluster generation with MiSeq, the sequencing files were processed and genomic sequence reads were obtained. The full genome sequence was assembled by CLC platform.

#### Phylogenetic analysis

Alignment of each segment was performed with the Mega (Version 6.05) software package; the phylogenetic trees of all the 8 segments were conducted by using neighbor-joining method with Kimura two-parameter distances. The bootstrap value was 1000.



hemagglutinin cleavage site of JX14737 possessed only one basic amino acid (PSRSSR↓GL) between HA1 and HA2, which indicated that this virus was a low pathogenic AIV in poultry (Table 1).

## Discussion

Subtype-H9N2 influenza virus was first identified in chicken farms in China in 1994 [19] and was found to be widespread in poultry and wild birds in Asia. Three major lineages of H9N2 virus circulated in poultry and wild birds: G1, Y280/G9, and BJ/94, represented by A/quail/Hong Kong/G1/97, A/duck/Hong Kong/Y280/97, and A/chicken/Beijing/1/1994 [20], respectively. H9N2 virus has been a gene donor for internal segments to many novel AIVs. Recently, many human infections with novel AIVs have occurred in China [3, 12, 21], especially in southern and eastern China. For instance, human infections with novel H7N9 and H10N8 AIVs were first reported in eastern China in 2013. Analysis of the full genomes of H7N9 and H10N8 viruses isolated from the patients showed that all 6 internal segments of these virus had originated in H9N2 AIVs [2, 4]. Since 2015, recent human infections with the high pathogenic avian influenza virus H5N6 have also been reported to have acquired internal genes from H9N2, although the first H5N6 virus found in humans had internal genes from H5N1 [22].

Sporadic human infections with H9N2 virus have been reported since the 1990s. Epidemiology data showed that some of these patients did not have direct contact with live poultry or any history of live poultry exposure [3]. These studies suggested that humans could be infected by AIVs without contacting live poultry directly. In this study, monthly air sample surveillance for AIV was conducted at 46 sites in 10 LPMs in Nanchang between April 2014 and March 2015. Of 807 air samples, 275 samples (34.1%) were Flu A positive by RRT-PCR, indicating a relatively high proportion of nucleic acid contamination in the air of LPMs. In addition, one (0.124%)

**Table 1** Key amino-acid changes in the full genome of JX14737 virus isolated from air samples

Protein	Amino-Acid Mutations	Phenotype
PB2	292 V	Increases polymerase activity in 293 T cells
	588 V	Enhanced polymerase activity and viral replication in human NHBE cells
HA	Q226L <sup>a</sup>	Increased viral binding to α2,6-sialic acid receptor, airborne transmission between mammals
	N/K137A <sup>a</sup>	Avian-like receptor binding preference (N/137 K)

<sup>a</sup>The amino acid positions of HA gene represent H3 numbering

H9N2 virus was isolated in the Jingdong bird and flower market, a large wholesale market in Nanchang city, Jiangxi province. The virus isolated from air samples was a low pathogenic AIV, but some mammalian-adaptation mutations were detected, along with mutations relating to polymerase activity and virus replication. Airborne virus has also been detected in LPMs in Guangdong [9] and Hong Kong [12] in South China, both of which are in or adjacent to tropical areas, while in this study it was found in Jiangxi province, a typical subtropical area in central China.

There were some limitations to this study. First, the air sample collection time (1.5 h) and the distance between stall center and instrument might affect the results. Second, the presence of polluted particles in the air might decrease the sensitivity rate of virus isolation.

## Conclusions

In our study, one H9N2 AIV was successfully isolated from 807 aerosolized samples. The virus shared high nucleotide homology with H9N2 AIVs from South China. Our study provided further evidence that the air in LPMs can be contaminated by influenza virus and its nucleic acids, and this fact should be considered when choosing and evaluating possible disinfection strategies for LPMs, such as regular air disinfection. Further statistical analysis suggested that the Flu A positive rate of air samples collected 3 m from poultry stalls were significantly higher than that of air samples collected at 5 and 7 m in the retail LPM ( $\chi^2 = 39.792$ ,  $p < 0.01$ ). The results suggested that the risk of human infection by aerosolized virus was increased when people were closer to stalls selling live poultry.

## Additional file

**Additional file 1: Figure S1.** Phylogenetic trees of the internal genes of JX14737. Trees were built by the neighbor-joining method using the MEGA6 software package (bootstrap value = 1000). The JX14737 H9N2 virus isolated from Jingdong Bird and Flower LPM in this study is highlighted by a red circle. The G57-like strain represented by A/chicken/Zhejiang/HJ/2007(H9N2) is highlighted by a blue square. The Y280-like strain represented by A/duck/Hong Kong/Y280/1997(H9N2) is highlighted by a pink rhombus. The G1-like strain represented by A/Quail/Hong Kong/G1/1997(H9N2) is highlighted by a green triangle. The phylogenetic trees of PB2, PB1, PA, NP, M, and NS genes of JX14737 are a-f, respectively. (PDF 343 kb)

## Abbreviations

AIVs: Avian Influenza Virus; GISAID: Global Initiative on Sharing All Influenza Data; LPMs: Live poultry markets; NGS: Next generation sequencing; RRT-PCR: Real-Time Reverse Transcription-Polymerase Chain Reaction

## Acknowledgements

We gratefully acknowledge the authors and laboratories who originated and submitted the sequences that we used in this manuscript to the GISAID EpiFlu database.

**Funding**

This research was supported by the National Key Research and Development Program of China (2016YFD0500208 to Wang and 2016YFC1200200 to Shu); National Natural Science Foundation of China (81,460,302 to H. Chen; 81,660,339 to M. Liu); The Major Science and Technology Project of Jiangxi Province (20143ACG70004 to H. Chen).

**Availability of data and materials**

The sequence of all eight segments of A/environment/Jiangxi/14737/2014 (H9N2) was submitted to the Global Initiative on Sharing All Influenza Data (GISAID), and the accession numbers were from EPI858159 to EPI858166.

**Authors' contributions**

SYL, CHY and WDY designed the study; ZXX, LMB, ZH, WJW, ZX, HFL, and FGY performed the experiments; CWB, YL, ZXX analyzed the data; and ZXX wrote the paper. All authors read and approved the final manuscript.

**Ethics approval and consent to participate**

Not applicable.

**Consent for publication**

Not applicable.

**Competing interests**

The authors declare that they have no competing interests.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 28 February 2017 Accepted: 10 July 2017

Published online: 24 July 2017

**References**

- Zhang Z, et al. The complexity of human infected AIV H5N6 isolated from China. *BMC Infect Dis.* 2016;16(1):600.
- Chen H, et al. Clinical and epidemiological characteristics of a fatal case of avian influenza A H10N8 virus infection: a descriptive study. *Lancet.* 2014; 383(9918):714–21.
- Butt KM, et al. Human infection with an avian H9N2 influenza A virus in Hong Kong in 2003. *J Clin Microbiol.* 2005;43(11):5760–7.
- Gao R, et al. Human infection with a novel avian-origin influenza A (H7N9) virus. *N Engl J Med.* 2013;368(20):1888–97.
- Yuan J, et al. Origin and molecular characteristics of a novel 2013 avian influenza A(H6N1) virus causing human infection in Taiwan. *Clin Infect Dis.* 2013;57(9):1367–8.
- Zhang R, et al. Clinical, epidemiological and virological characteristics of the first detected human case of avian influenza A(H5N6) virus. *Infect Genet Evol.* 2016;40:236–42.
- Yuan R, et al. Continuing Reassortant of H5N6 subtype highly pathogenic avian influenza virus in Guangdong. *Front Microbiol.* 2016;7:520.
- Chen LJ, et al. Diversity and evolution of avian influenza viruses in live poultry markets, free-range poultry and wild wetland birds in China. *J Gen Virol.* 2016;97(4):844–54.
- Wu Y, et al. Aerosolized avian influenza A (H5N6) virus isolated from a live poultry market, China. *J Inf Secur.* 2017;74(1):89–91.
- Zhou J, et al. Isolation of H5N6, H7N9 and H9N2 avian influenza A viruses from air sampled at live poultry markets in China, 2014 and 2015. *Euro Surveill.* 2016;21(35).
- Zhong L, et al. Molecular mechanism of the airborne transmissibility of H9N2 avian influenza A viruses in chickens. *J Virol.* 2014;88(17):9568–78.
- Sun Y, Liu J. H9N2 influenza virus in China: a cause of concern. *Protein Cell.* 2015;6(1):18–25.
- Guo Y, Xie J, Wu K, Dong J, et al. Characterization of genome of a/ Guangzhou/333/99(H9N2) virus. *Chin J Exp Clin Virol.* 2002;16:142–5.
- Zhou B, Donnelly ME, Scholes DT, et al. Single-reaction genomic amplification accelerates sequencing and vaccine production for classical and swine origin human influenza A viruses. *J Virol.* 2009;83(19):10309–13.
- Rutvisuttinunt W, et al. Simultaneous and complete genome sequencing of influenza A and B with high coverage by Illumina MiSeq platform. *J Virol Methods.* 2013;193(2):394–404.
- Wan H, Perez DR. Amino acid 226 in the Hemagglutinin of H9N2 influenza viruses determines cell tropism and replication in human airway epithelial cells. *J Virol.* 2007;81(10):5181–91.
- Liu J, Wu Q, Chen F, et al. Comparison of amino acids in the receptor-binding site in hemagglutinin between chicken and human H9N2 influenza virus. *Chin J Prev Vet Med.* 2003;25(6):416–8.
- Xiao C, et al. PB2-588 V promotes the mammalian adaptation of H10N8, H7N9 and H9N2 avian influenza viruses. *Sci Rep.* 2016;6:19474.
- Subbarao K, et al. Genetics, receptor binding property, and transmissibility in mammals of naturally isolated H9N2 avian influenza viruses. *PLoS Pathog.* 2014;10(11):e1004508.
- Guo YJ, et al. Characterization of the pathogenicity of members of the newly established H9N2 influenza virus lineages in Asia. *Virology.* 2000; 267(2):279–88.
- Guo Y, Li J, Cheng X, et al. Discovery of men infected by avian influenza A(H9N2) virus. *Chin J Exp Clin Virol.* 1999;13(2):1–6.
- Xu W, et al. A fatal case of infection with a further reassortant, highly pathogenic avian influenza (HPAI) H5N6 virus in Yunnan, China. *Infect Genet Evol.* 2016;40:63–6.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
www.biomedcentral.com/submit

